



Large-scale displacement along the Altyn Tagh Fault (North Tibet) since its Eocene initiation: Insight from detrital zircon U–Pb geochronology and subsurface data

Feng Cheng, Marc Jolivet, Suotang Fu, Changhao Zhang, Qiquan Zhang, Zhaojie Guo

► To cite this version:

Feng Cheng, Marc Jolivet, Suotang Fu, Changhao Zhang, Qiquan Zhang, et al.. Large-scale displacement along the Altyn Tagh Fault (North Tibet) since its Eocene initiation: Insight from detrital zircon U–Pb geochronology and subsurface data. *Tectonophysics*, 2016, 677-678, pp.261-279. 10.1016/j.tecto.2016.04.023 . insu-01312745

HAL Id: insu-01312745

<https://hal-insu.archives-ouvertes.fr/insu-01312745>

Submitted on 20 May 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Large-scale displacement along the Altyn Tagh Fault (North Tibet) since its
Eocene initiation: insight from detrital zircon U-Pb geochronology and
subsurface data**

Feng Cheng ^{1,2}, Marc Jolivet ², Suotang Fu ³, Changhao Zhang ³, Qiquan Zhang ³,
Zhaojie Guo ^{1,†}

¹ *Key Laboratory of Orogenic Belts and Crustal Evolution, Ministry of Education,
School of Earth and Space Sciences, Peking University, Beijing, 100871, China*

² *Laboratoire Géosciences Rennes, CNRS-UMR6118, Université Rennes 1 -
Observatoire des Sciences de l'Univers, Rennes, 35042, France*

³ *Qinghai Oilfield Company, PetroChina, Dunhuang, Gansu, 736202, China*

[†] *Corresponding author. Tel.: + 86-10-62753545; fax: + 86-10-62758610. E-mail
address: zjguo@pku.edu.cn (Z.J. Guo)*

16 **Abstract**

17 Marking the northern boundary of the Tibetan plateau, the Altyn Tagh fault plays
18 a crucial role in accommodating the Cenozoic crustal deformation affecting the
19 plateau. However, its initiation time and amount of offset are still controversial
20 despite being key information for the understanding of Tibet evolution. In this study,
21 we present 1122 single LA-ICP-MS detrital zircon U-Pb ages obtained from 11
22 Mesozoic to Cenozoic sandstone samples, collected along two sections in the
23 northwestern Qaidam basin (Eboliang and Huatugou). These data are combined with
24 new 3D seismic reflection profiles to demonstrate that: (1) From the Paleocene to
25 early Eocene, the Eboliang section was approximately located near the present
26 position of Anxi, 360 ± 40 km southwest from its current location along the Altyn
27 Tagh fault, and sediments were mainly derived from the Altyn Tagh Range. At the
28 same period, the Huatugou section was approximately located near the present
29 position of Tula, ca. 360 km southwest from its current location along the Altyn Tagh
30 fault, and the Eastern Kunlun Range represented a significant sediment source. (2)
31 Left-lateral strike-slip movement along the Altyn Tagh fault initiated during the
32 early-middle Eocene, resulting in northeastward displacement of the two sections. (3)
33 By early Miocene, the intensive deformation within the Altyn Tagh Range and
34 northwestern Qaidam basin strongly modified the drainage system, preventing the
35 materials derived from the Altyn Tagh Range to reach the Eboliang and the Huatugou
36 sections. The post-Oligocene clastic material in the western Qaidam basin is generally

derived from local sources and recycling of the deformed Paleocene to Oligocene strata. From these data, we suggest enhanced tectonic activity within the Altyn Tagh Range and northwestern Qaidam basin since Miocene time, and propose an early-middle Eocene initiation of left-lateral strike-slip faulting leading to a 360 ± 40 km offset along the Altyn Tagh fault.

Keywords: Detrital zircon U-Pb geochronology, Cenozoic tectonics, North Tibet, Altyn Tagh fault, Qaidam basin.

1. Introduction

Two competing end-member mechanisms have been proposed to explain the accommodation of the ongoing convergence between India and Eurasia since the early Eocene collision: (1) a homogeneous crustal thickening of the Tibetan plateau (e.g. England and Houseman, 1989; Searle, 1996); and (2) an eastward extrusion of the Tibetan plateau and southeast Asia away from the indenting Indian plate (Molnar and Tapponnier, 1975; Peltzer and Tapponnier, 1988; Avouac and Tapponnier, 1993; Tapponnier, et al., 2001). The second model requires large-scale displacement along lithospheric strike-slip fault zones to allow extrusion of the Tibetan crust (e.g. Searle, 1996; Tapponnier, et al., 2001). Marking the northern boundary of the Tibetan plateau (Fig. 1), the lithospheric-scale left-lateral strike-slip Altyn Tagh fault (ATF) plays a crucial role in accommodating the crustal deformation and appears to be an ideal field laboratory for ascertaining the dynamics of plateau formation (Molnar and

Tapponnier, 1975; Wittlinger et al., 1998; Jolivet et al., 1999, 2001; Yin and Harrison, 2000; Yin et al., 2002; Searle et al., 2011). Understanding the kinematic pattern of the ATF, especially the exact Cenozoic initiation time and the amount of Cenozoic offset, is of major importance for unraveling the crustal accommodation processes within the plateau since the India-Eurasian collision and for deciphering the growth history of the entire Tibetan plateau.

Several authors proposed that a large scale Jurassic basement cooling event associated to tectonic exhumation affected a corridor along the ATF (e.g. Delville et al., 2001; Sobel et al., 2001; Wang et al., 2005). In addition, many researchers consider that deformation along the ATF initiated after the late Mesozoic, during the early Cenozoic, or even during the Neogene (e.g. Tapponnier et al., 1986; Jolivet et al., 1999; Chen et al., 2001; Jolivet et al., 2001; Yin et al., 2002; Wang et al., 2006a; Wu et al., 2012a, 2012b; Zhang et al., 2012, 2014a). Similarly, estimations of the total displacement along the ATF vary widely from ~1200 km to less than 90 km (e.g. Tapponnier et al., 1986; Wang, 1997; CSBS, 1992; Ritts and Biffi, 2000; Yin and Harrison, 2000; Yang et al., 2001; Yue et al., 2001; Chen et al., 2002; Yin et al., 2002; Cowgill et al., 2003; Gehrels et al., 2003a, 2003b; Darby et al., 2005; Yue et al., 2005; Searle et al., 2011; Cheng et al., 2015a). These tremendous discrepancies may be partially attributed to the immense size and extent of the Altyn Tagh Range, making it difficult to locate ideal piercing points to estimate the total displacement along the ATF. Furthermore, due to strong Cenozoic deformation, continuous Mesozoic to Cenozoic stratigraphic sections necessary to estimate the time of initiation of

left-lateral slip movement are seldom preserved (e.g. Yin and Harrison, 2000; Cheng et al., 2015a).

Detrital zircon analysis of a continuous, well dated stratigraphic succession has become a powerful tool for unraveling source to sink relationships and constraining the tectonic and topographic evolution of an area (e.g. Fedo et al., 2003; Thomas, 2011; Gehrels, 2014). To bring more constraints on the kinematic evolution of the ATF, we conducted an integrated analysis on two Jurassic to Pleistocene sedimentary sections in the western part of the Qaidam basin, adjacent to the ATF (Fig. 2). We then combined the detrital zircon U-Pb geochronology data obtained from these sections with high-quality subsurface data, including newly acquired seismic profiles and drill core sandstone samples.

2. Geological Background

2.1 Altyn Tagh Range

The Altyn Tagh Range is located along the northern edge of the Tibetan plateau, separating the Tarim basin to the northwest from the Tibetan plateau and the Qaidam basin to the south (Figs. 1 and 2). The bedrock of the Altyn Tagh Range mainly consists of Precambrian igneous and metamorphic rocks and Paleozoic igneous and sedimentary rocks (e.g. Sobel and Arnaud, 1999; Yin et al., 2002). It has been shown that the Altyn Tagh Range experienced multiple-stage deformation and tectonic exhumation from the Jurassic to the Holocene (Tapponnier et al., 1986; Jolivet et al., 1999; Yue and Liou, 1999; Chen et al., 2001; Delville et al., 2001; Jolivet et al., 2001;

Sobel et al., 2001; Yin et al., 2002; Wang et al., 2005, 2006a; Zhang et al., 2012). Within this range, the over 1600 km long ENE-trending ATF links the western Kunlun thrust belt to the southwest and the Qilian Shan thrust belt to the northeast (Fig. 1; Burchfiel et al., 1989; Wang, 1997; Yue and Liou, 1999; Yin and Harrison, 2000; Yin et al., 2002). Although Mesozoic shearing occurred in the Altyn Tagh Range, the growth of the northern Tibetan plateau is largely influenced by Cenozoic sinistral strike-slip faulting along the ATF (Tapponnier et al., 1986, 2001; Arnaud et al., 2003; Wang et al., 2005; Li et al., 2006; Liu et al., 2007). As previously mentioned, the estimation of the total displacement along the fault remains heavily debated, varying from about 1200 km to less than 90 km (e.g. CSBS, 1992; Ritts and Biffi, 2000; Yang et al., 2001; Chen et al., 2002; Cowgill et al., 2003; Gehrels et al., 2003a, 2003b; Yue et al., 2005; Cheng et al., 2015a).

2.2 *Qaidam basin*

The rhomb-shaped Qaidam basin is the largest petroliferous basin within the entire Tibetan plateau (Figs. 1 and 2). Geomorphologically, the basin is surrounded by the Altyn Tagh Range to the northwest, the Qilian Shan to the northeast and the Eastern Kunlun Range to the south. Geological mapping and petroleum exploration revealed that the Qaidam basin is filled with Mesozoic to Cenozoic clastic sediments unconformably overlying a poorly documented basement (Xia et al., 2001; Meng and Fang, 2008; Yin et al., 2008a, 2008b; Zhang et al., 2013a). The Mesozoic strata, especially the Jurassic and lower Cretaceous sequences, are mainly distributed along the foreland of both the Altyn Tagh Range and the Qilian Shan (Ritts et al., 1999; Wu

et al., 2011), while the Cenozoic series are generally deposited over the entire basin (Yin et al., 2008b). The Cenozoic sedimentation pattern is largely controlled by a succession of depocenters consistently located along the long axis of the basin and gradually migrating eastward since Eocene times, indicating the gradual uplift of the Altyn Tagh Range (Fig. 1; Song and Wang, 1993; Qiu, 2002; Sun et al., 2005; Wang et al., 2006a; Yin et al., 2008b). Using magnetostratigraphy, palynology and paleontology, the Mesozoic-Cenozoic strata have been precisely subdivided into 11 chronostratigraphically constrained units (Fig. 3; Huo, 1990; QBGMR, 1991; Yang et al., 1992; Huang et al., 1996; Xia et al., 2001; Qiu, 2002; Deng et al., 2004a, 2004b; Sun et al., 2005; Zhang, 2006; Zhao et al., 2006; Fang et al., 2007; Sun et al., 2007; Yin et al., 2008b; Gao et al., 2009; Pei et al., 2009; Lu and Xiong, 2009; Ke et al., 2013). These units are: (1) the Dameigou Formation, $J_{1+2}d$; (2) the Caishiling Formation, J_3c ; (3) the Quanyagou Formation, Kq ; (4) the Lulehe Formation, $E_{1+2}l$, >53.5 - 43.8 Ma (Yang et al., 1992; Zhang, 2006; Ke et al., 2013); (5) the lower Xiaganchaigou Formation, E_3^1xg , 43.8 - 37.8 Ma (Zhang, 2006; Sun et al., 2007; Pei et al., 2009); (6) the upper Xiaganchaigou Formation, E_3^2xg , 37.8 - 35.5 Ma (Sun et al., 2005; Sun, 2007; Pei et al., 2009); (7) the Shangganchaigou Formation, N_1sg , 35.5 - 22.0 Ma (Sun et al., 2005; Lu and Xiong, 2009); (8) the Xiayoushashan Formation, N_2^1xy , 22.0 - 15.3 Ma (Fang et al., 2007; Lu and Xiong, 2009); (9) the Shangyoushashan Formation, N_2^2sy , 15.3 - 8.1 Ma (Fang et al., 2007); (10) the Shizigou Formation, N_2^3s , 8.1 - 2.5 Ma (Fang et al., 2007); (11) the Quaternary deposits, including the Qigequan Formation (Q_1q) and the Dabuxun-Yanqiao

Formation (Q_2d), 2.5 - 0.01 Ma (Fang et al., 2007; Yin et al., 2008b). Effective elastic thickness calculation implies that the mechanical strength of the Qaidam crust is exceptionally strong compared to the rest of the Tibetan plateau (Braitenberg et al., 2003). Based on balanced cross section results, Zhou et al. (2006) proposed that the Qaidam basin experienced an average of 10% of NE-SW shortening during the Cenozoic, whereas Yin et al. (2008b) suggested that the shortening strain across the basin decreases systematically eastward from ca. 48% in the west, to ca. 11% in the center, and <1% in the east.

3. Stratigraphy and sedimentary characteristics of the studied sections

In this study, the Eboliang and Huatugou sections, situated along the eastern and central segments of the ATF, have been chosen for their exceptional preservation and exposure of the Jurassic to Pleistocene sedimentary series (Figs. 2 and 3). In order to give a continuous description of the Jurassic to Quaternary sediment evolution of the northwestern Qaidam basin, we constructed the Eboliang and Huatugou sections based on the lithologies, sediment facies and paleocurrent directions obtained from fieldwork, drill-core and literature data as summarized in the Figure 3 and in the text below. The stratigraphy of the Huatugou and Eboliang sections has already been established by previous field geological studies which give age constraints based on fossil assemblages, distinctive lithology feature, magnetostratigraphy, palynology and paleontology (e.g. XBGRM, 1993; HGSI, 2003; IGSQP, 2004). Consequently, we follow the previous division of formations.

3.1 Eboliang section

The Eboliang section is located along the southern flank of the Altyn Tagh Range (Fig. 2). The upper Jurassic-Cretaceous strata are missing, and the Paleogene deposits rest unconformably on lower - middle Jurassic series. Except for the absence of the Paleocene Lulehe Formation, the Cenozoic sequence is complete and exceptionally well exposed.

High-quality seismic profiles and drill core data reveal that the base of the Eboliang section is formed by the lower to middle Jurassic Dameigou Formation (J_{1+2d}), resting unconformably on the basement rocks (Yin et al., 2008). This Jurassic sequence generally contains a succession of three well-developed fining upward cycles. Each cycle characteristically begins with medium- to coarse-grained sandstone deposits (Fig. 4A), fining upwards to silty mudstone. The sequence, as in general for the Qaidam basin, is considered to represent shore shallow lacustrine facies at the base, evolving into flood plain sediments upward (Ritts et al., 1999; Ritts and Biffi, 2000; Wu et al., 2011; Jian et al., 2013). Trough and planar cross-stratification and clast imbrication from those Jurassic fluvial strata indicates northward paleoflows (Fig. 3; Ritts and Biffi, 2000).

The entire upper Jurassic and Cretaceous sequence is missing and the Paleocene Lulehe Formation (E_{1+2l}) unconformably overlies the lower and middle Jurassic series (Wu et al., 2011). The strata are composed of dark brown mudstone (Fig. 4B) and sandy mudstone intercalated with gray siltstone at the base, evolving towards brown pebbly sandstone intercalated with conglomerates and gypsum-salt layers in

the middle and upper parts of the section. The Lulehe Formation is generally interpreted as braided river and alluvial fan depositional environments (Zhuang et al., 2011; Jian et al., 2013; Song et al., 2013). Based on heavy mineral contents, calculated zircon-tourmaline-rutile (ZTR) indices, and the heavy mineral assemblages from drill-core samples from a large number of oil wells in the Eboliang section, Fu et al. (2013) suggested that during the deposition of the Paleocene-early Eocene Lulehe Formation, the material deposited in the Eboliang area was derived from the north, probably from the Altyn Tagh Range.

The Eocene to Oligocene strata of the Xiaganchaigou Formation unconformably rest on the Lulehe Formation. The strata are mainly composed of grey and brown sandstone (Fig. 4C) and conglomerate at the bottom corresponding to alluvial facies deposits (Zhuang et al., 2011; Jian et al., 2013; Wang et al., 2013) and contain a succession of fining upward cycles. The middle part of the Xiaganchaigou Formation is dominated by brownish red mudstone and sandy mudstone intercalated with gray calcareous mudstone and marlstone. The upper part of the Xiaganchaigou Formation, however, is dominated by variegated mudstone intercalated with thin-layers of sandy conglomerate, sandstone and limestone, generally associated to a lacustrine depositional environment (Zhuang et al., 2011; Jian et al., 2013; Wang et al., 2013). Conglomerate fabrics in the Eboliang area show north-directed or south-directed bilateral paleoflows, suggesting a low-energy lacustrine-offshore environment at the time (Fig. 3; Wu et al., 2012b).

The Oligocene Shangganchaigou Formation (N₁sg), conformably rests on the

210 Xiaganchaigou Formation. It consists of brownish red, gray siltstone, grey mudstone,
211 sandy mudstone and siltstone, intercalated with gray sandstone and marlstone (Fig.
212 4D) mostly corresponding to lacustrine sediment facies (Zhuang et al., 2011; Jian et
213 al., 2013; Wang et al., 2013). Paleocurrents measured from the conglomerate fabrics
214 become dominantly south-directed away from the Altyn Tagh Range (Fig. 3; Wu et al.,
215 2012b).

216 The early Miocene Xiayoushashan Formation (N_2^1xy) conformably lies on the
217 Shangganchaigou Formation. The lower part of the formation is composed of gray
218 and brownish red mudstone and sandy mudstone intercalated with gray siltstone,
219 grayish yellow sandstone and grayish brown marlstone. It evolves upwards to gray
220 mudstone, argillaceous sandstone and argillaceous siltstone intercalated with grey
221 siltstone, brownish red mudstone and gray marlstone (Fig. 4E). The early Miocene
222 sequence in the north Qaidam basin is generally considered as deposited in a fluvial to
223 marginal lacustrine environment (Zhuang et al., 2011; Jian et al., 2013; Wang et al.,
224 2013). Conglomerate fabrics in the section suggest south-directed paleocurrents
225 during the early Miocene (Fig. 3; Wu et al., 2012b).

226 The late Miocene Shangyoushashan Formation (N_2^2sy) conformably rests on the
227 Xiayoushashan Formation. The deposits are mainly composed of grey mudstone and
228 sandy mudstone intercalated with grey argillaceous sandstone, argillaceous siltstone,
229 grayish yellow marlstone and some limited gray sandy limestone corresponding again
230 to a fluvial to marginal lacustrine environment (Zhuang et al., 2011; Jian et al., 2013;
231 Wang et al., 2013). Paleocurrent measurements from the sandstone layers around the

Eboliang area indicate southwest-directed paleoflows (Fig. 3; Wu et al., 2012b).

The Pliocene Shizigou Formation (N_2^3s) generally conformably and locally unconformably rests on the Shangyoushashan Formation. The series are composed of gray mudstone and silty mudstone intercalated with brown siltstone, pebbly sandstone and gypsum-salt layers. They are generally interpreted as deposited in a marginal lacustrine to alluvial fan environment (Fig. 4F; Zhuang et al., 2011; Heermance et al., 2013; Jian et al., 2013; Wang et al., 2013). Trough cross-bedding within the strata suggests generally southwestward paleoflows during the Pliocene (Fig. 3; Heermance et al., 2013).

Finally, the Pleistocene sequence, dominated by the Qigequan Formation (Q_{1q}), unconformably overlies the Shizigou Formation. The sediments are characterized by brown sandstone, pebbly sandstone and conglomerate intercalated with gypsum layers, corresponding to fluvial to evaporitic lacustrine facies (Heermance et al., 2013; Jian et al., 2013). Well-developed cross-beddings within the strata show primarily southward unidirectional paleocurrents (Fig. 3; Heermance et al., 2013).

3.2 Huatugou section

The Huatugou section is located in the northernmost part of the prominent Yingxiongling uplift structure (Fig. 2). Except again for the Paleocene Lulehe Formation, the whole Mesozoic-Cenozoic sequence is well exposed (Figs. 2 and 3).

Subsurface data reveal that the Jurassic clastic coal-bearing strata rest unconformably on the Altyn Tagh Range basement (Xia et al., 2001; Yin et al., 2008).

The Dameigou Formation ($J_{1+2}d$) is dominated by a set of sandstone beds interbedded

with conglomerate, siltstone and mudstone (Ritts et al., 1999; Ritts and Biffi, 2000). The sediment facies vary from fluvial to shallow lake deposits (Ritts and Biffi, 2000). Trough and planar cross-stratification as well as clast imbrication within the strata indicate northward paleoflows during the Jurassic (Fig. 3; Ritts and Biffi, 2000).

Based on the subsurface data (core and seismic profile data), the Paleocene to early Eocene Lulehe Formation (E_{1+2l}) is dominated by successions of purple-red conglomerate interbedded with sandstone at the base, evolving upward towards grey sandstone, siltstone and sandy mudstone. The sediment facies vary from fluvial to shallow lake deposits upward and basinward (Fu et al., 2013; Li et al., 2015). Heavy mineral contents in surface samples, calculated zircon-tourmaline-rutile (ZTR) indices, and the heavy mineral assemblages from the drill-core samples from a large number of oil wells in the Huatugou area suggest that during the deposition of the Paleocene-early Eocene Lulehe Formation, the material deposited in that area was mainly derived from the southwest, probably from the Eastern Kunlun Range (Fu et al., 2013; Li et al., 2015).

The middle Eocene lower Xiaganchaigou Formation (E_3^1xg) is mainly composed of grey-green, brownish-red sandstone and pebbly sandstone at the base, evolving upward towards brown red sandstone and siltstone (Fig. 4G). The sediment facies vary from fluvial to shallow lacustrine deposits upward (Zhuang et al., 2011; Wu et al., 2012a, 2012b). Paleocurrents measurements based on pebble imbrication and cross-stratification in the fluvial sandstone layers indicate NE-directed paleoflows (Fig. 3; Meng and Fang, 2008).

The late Eocene upper Xiaganchaigou Formation (E_3^{2xg}) rests conformably on the lower Xiaganchaigou Formation. This sequence is dominated by successions of grey-green, brownish-red mudstone and carbonaceous siltstone, intercalated with argillaceous siltstone and marlstone (Fig. 4H). The sediment facies vary upward from braided river to shallow lake deposits (Wu et al., 2012a, 2012b). Clast imbrications within the fluvial strata suggest that paleocurrents were directed towards the south during the deposition of the upper Xiaganchaigou Formation. (Fig. 3; Wu et al., 2012a).

The Oligocene Shangganchaigou Formation (N_1sg), conformably resting on the upper Xiaganchaigou Formation, is mainly composed of greyish-green carbonaceous mudstone and sandstone interbedded with grey-white mudstone. This series is dominated by deltaic to shallow lake facies deposits (Wu et al., 2012a, 2012b). Reported paleocurrents from clast imbrications are generally directed towards the south (Fig. 3; Wu et al., 2012a).

The Xiayoushashan Formation (N_2^1xy), conformable with the Shangganchaigou Formation, is mainly composed of yellowish-brown, pebbly sandstone, sandstone and argillaceous siltstone with intercalations of limestone. The facies vary from shallow lake to fluvial deposits (Wu et al., 2012a, 2012b). Clast imbrications within the sequences suggest generally southwestward paleoflows during the early Miocene.

The Shangyoushashan Formation (N_2^{2sy}), resting unconformably on the Xiayoushashan Formation (Wang et al., 2010), mainly consists of yellow pebbly sandstone, sandstone and silty mudstone. The sediment facies is dominated by

shallow lake, alluvial plain and braided river deposits (Wu et al., 2012a, 2012b). Clast imbrication as well as trough (and lesser planar) cross-stratification within the layers in the section suggest generally northwestward paleoflows (Zhuang et al., 2011).

The Shizigou Formation (N_2^3s), again lying unconformably on the Shangyoushashan Formation, is mainly composed of earth-yellow sandstone and silty mudstone corresponding to fluvial facies sediments (Wu et al., 2012a, 2012b). Paleocurrents from clast imbrications indicate generally south-directed paleoflows away from the Altyn Tagh Range during the deposition of the Shizigou Formation (Fig. 3; Zhuang et al., 2011).

The Qigequan (Q_1q) and Dabuxun-Yanqiao (Q_2) formations, unconformably overly the Shizigou Formation, and mainly consists of grey and brown sandstone and conglomerate corresponding to flood plain and alluvial facies deposits.

4. Methods and analytical procedures

4.1 Detrital zircon geochronology

Detrital zircon U-Pb geochronology has rapidly developed into a very powerful tool for determining sediment provenances (e.g. Fedo, et al., 2003; Thomas, 2011; Liu et al., 2013; Yang et al., 2013; Gehrels et al., 2014; Cheng et al., 2015a). By systematically comparing the detrital zircon U-Pb age spectrum obtained from sedimentary sequences in basins with the known ages of potential source terranes, it is possible to describe the source to sink relations through time within a given area and to reconstruct the landscape evolution of the region (e.g. Fedo, et al., 2003; Gehrels et

al., 2011; Thomas, 2011; Liu et al., 2013; Yang et al., 2013; Gehrels et al., 2014; Yang et al., 2014; Cheng et al., 2015a). In the Eboliang section, eight samples were collected, ranging in age from Jurassic to Pleistocene (Fig. 2), and including three samples from drill cores. In the Huatugou section, three core samples were obtained from drill wells, ranging in age from Paleocene to Oligocene. In order to derive a continuous Jurassic to Pleistocene source to sink relation between the Altyn Tagh Range and the western Qaidam basin, we also integrated six published detrital zircon U-Pb dating results obtained on samples collected from the Huatugou section (Cheng et al., 2015a, 2016). The major petrological characteristics of all the samples are described in Table 1.

Zircon grains for U-Pb age dating were concentrated from each sample following the standard procedures outlined in Liu et al. (2013). This work was conducted at the Chengxin Geology Service Co. Ltd, Langfang, Hebei Province, China. Individual zircon crystals (generally more than 200 grains) were mounted in epoxy resin without handpicking to avoid sampling bias. Samples were then polished to obtain a smooth flat internal surface. Reflected and transmitted light as well as cathodoluminescence (CL) images were made to reveal internal heterogeneities and allow choosing potential internal targets for isotopic dating. U-Pb analysis was performed on an Agilent 7500a ICP-MS connected to an American New Wave UP 193 SS 193 nm Excimer laser ablation system at the China University of Geosciences, Beijing. All samples were analyzed using a laser spot size of 36 μm and a frequency of 10 Hz. Two standards (Black et al., 2003; Qi et al., 2005) were analyzed every 10 to 20

grains, to correct for instrument fluctuations and determine fractionation factors. Zircons Qinghu and 91500 (Wiedenbeck et al., 1995) were the monitoring standards. For elemental concentration analysis, NIST610 was the external standard, and ^{29}Si was the internal standard. Meanwhile, NIST612 and NIST614 were used as monitoring standards. The GLITTER 4.4 software was used to calculate the U-Pb isotope ratios and element contents (China University of Geosciences, Lab. of Prof. Y.S. Liu, Beijing, China). The U-Pb ages obtained were checked for discordance by plotting the analyses on concordia diagrams using the Isoplot 3.0 software (Ludwig, 2003). The common-Pb correction followed the method described by Andersen (2002). Ages younger than ca. 1000 Ma are based on common Pb-corrected $^{206}\text{Pb}/^{238}\text{U}$ ratios, whereas ages older than ca. 1000 Ma are based on common Pb-corrected $^{207}\text{Pb}/^{206}\text{Pb}$ ratios. For ICP-MS analyses, those ages with discordance degree >10% were excluded from analysis (e.g. Gehrels et al., 2003a; Yang et al., 2013; Liu et al., 2015). A more complete description of the sample separation methods and analytical procedures is given in Yuan et al. (2004).

4.2 Seismic profile

Extensive petroleum exploration of the Qaidam basin in recent years has provided abundant subsurface data, including high-quality seismic profiles and drill core data. In this study, we integrated two 3D seismic blocks and two 2D seismic profiles (A-A', B-B', CC' and DD', see locations in Figs. 1 and 2) with our surface field investigation to describe the tectonic history of the western Qaidam basin. Seismic data were interpreted using the SMT Kingdom software.

5. U–Pb geochronology results of detrital zircons

In general, about 90% of the zircon crystals are characterized by relatively distinct oscillatory zoning in CL images and relative high Th/U ratios, indicating a magmatic origin (Corfu et al., 2003; Hanchar and Rudnick, 1995; Hoskin and Black, 2000). The detrital zircons U-Pb ages vary widely between 2956 Ma and 57. Except for a single Cenozoic U-Pb age in sample EBLE, the detrital zircons U-Pb ages can be statistically subdivided into three groups of Precambrian (spanning from ca. 2.8 Ga to 550 Ma), early to middle Paleozoic (peaks at 460~400 Ma) and late Paleozoic to Mesozoic (peaks at ca. 260~240 Ma). Representative CL images of typical zircon grains are presented in Figure 5. U-Pb isotopic ages with errors and related raw data are listed in full in Appendix A. The statistical U–Pb geochronology data for each of the samples are listed in Table 1 and a detailed description of the zircons analyzed in each sample is given in Appendix B. Concordia plots for the eleven samples are shown in Figure 6. The zircon age spectrum of the samples from the Eboliang and Huatugou sections are shown in Figures 7 and 8, respectively. In the following, major age groups and their corresponding peak ages have been established from visual inspection of the detrital zircon U-Pb age probability plots for all the samples. Age peaks are considered major when including at least 20% of the total number of data spread over less than 250 Ma, whereas a minor peak refers to populations representing less than 20% of the total number of data distributed over more than 300 Ma.

6. Discussion

6.1 Geochronological characteristics of potential sources for the sedimentary rocks in the western Qaidam basin

The Altyn Tagh Range, the Qilian Shan, as well as the Eastern Kunlun Range are all potential source regions likely to provide zircons to the Qaidam basin (e.g. Métivier et al., 1998; Xia et al., 2001; Meng and Fang, 2008; Yin et al., 2007, 2008a, 2008b). In order to better constrain the provenance area of the samples collected in the western Qaidam basin (Fig. 9A), we compiled the zircon U-Pb ages available on basement rocks of the Altyn Tagh Range (Fig. 9B), the Qilian Shan (Fig. 9C) and the Eastern Kunlun Range (Fig. 8D).

6.1.1 Altyn Tagh Range

In the Altyn Tagh Range, the basement mainly consists of Archean and Proterozoic rocks with Archean zircon U-Pb ages ranging from ~3.6 Ga to ~2.6 Ga (Lu and Yuan, 2003; Lu et al., 2008; Long et al., 2014; Zhang et al., 2014b and references therein), and Proterozoic zircon U-Pb ages ranging from ~2.4 Ga to ~650 Ma (Gehrels et al., 2003a, 2003b; Wang et al., 2006b; Lu et al., 2008 and references therein; Zhang et al., 2011; Wang et al., 2013). A few scattered Neoproterozoic intrusions are exposed, with a large, distinctive early Neoproterozoic intrusion (ca. 850 Ma to ca. 1000 Ma in age) exposed west of the Xorkol basin (Fig. 9A). Wang et al. (2013) recently reported a mean crystallization age of ~910 Ma for the basement rocks near the Anxi area, in the western part of the Altyn Tagh Range (Fig. 9A).

Paleozoic intrusions with ages spanning from ~550 Ma to ~400 Ma are widely distributed (Jolivet et al., 1999; Sobel and Arnaud, 1999; Zhang et al., 2001; Cowgill et al., 2003; Gehrels et al., 2003a; Chen et al., 2004; Yue et al., 2004a, 2005; Yang et al., 2006; Wang et al., 2014b and references therein). Finally a few Permian igneous rocks are exposed along the central part of the Altyn Tagh Range, with zircon U-Pb ages ranging between ~300 Ma and ~260 Ma (Fig. 9A; Cowgill et al., 2003; Gehrels et al., 2003a; Gehrels et al., 2003b; Wu et al., 2014).

6.1.2 *Qilian Shan*

Zircon U-Pb ages from the North Qaidam and South Qilian Shan terranes mainly group between 2700 to 1100 Ma, 550 to 400 Ma, and 300 to 200 Ma (Fig. 9B; Yang and Song, 2002; Cowgill et al., 2003; Gehrels et al., 2003a, 2003b; Yue et al., 2005; Shi et al., 2006; Song et al., 2014 and references therein). The North Qilian Shan basement mainly consists of early Paleozoic marine strata associated with a series of Ordovician volcanics and Silurian granitic plutons (Bovet et al., 2009; Xiao et al., 2009; Song et al., 2014 and references therein). These granitoid intrusions yielded U-Pb zircon ages ranging from ca. 550 to ca. 440 Ma (Wu et al., 2001; Gehrels et al., 2003a, 2003b; Yue et al., 2005; Song et al., 2014 and references therein). Finally, the central Qilian Shan basement is composed of Mesoproterozoic to Neoproterozoic marine sequences intruded by early Paleozoic plutons, with detrital and plutonic zircon ages comprised between 2333 to 874 Ma and 442 to 424 Ma respectively (Guo and Li, 1999; Lu, 2002; Yue et al., 2005; Bovet et al., 2009; Song et al., 2014 and references therein). The Qilian Shan basement was affected by a Devonian phase of

metamorphism (400-360 Ma), with a metamorphic peak at 440-423 Ma within the north Qaidam ultrahigh pressure belt (Yang et al., 2001, Yang and Song, 2002; Song et al., 2006; Yang et al., 2006; Song et al., 2014 and references therein). Finally, sporadically distributed late Permian to early Triassic granitoids are exposed along the western edge of the South Qilian Shan terrane, yielding zircon U-Pb ages ranging from ca. 270 Ma to ca. 230 Ma (Fig. 9A and 9C; Yang and Song, 2002; Wu et al., 2009; Dong et al., 2014, 2015).

6.1.3 Eastern Kunlun Range

Within the Eastern Kunlun Range, Proterozoic ages are rare (Fig. 9A and 9D). Several early Paleozoic intrusions with U-Pb zircon ages ranging from ca. 500 Ma to 400 Ma have been reported by previous studies (Fig. 9C; Cowgill et al., 2003; IGSQP, 2004; Dai et al., 2013; Li et al. 2013). Late Paleozoic - early Mesozoic granitoids (ca. 300 Ma to ca. 200 Ma), corresponding to the magmatism associated to the Permo-Triassic closure of the Paleo-Tethys Ocean and to the post-collision magmatism that followed the docking of the Qiangtang Block, are extensively distributed (Fig. 9A; e.g. Roger et al., 2003, 2008, 2010; Liu et al., 2006; Li et al., 2013; Jolivet et al., 2015; Chen et al., 2015). Finally, a few Cretaceous ages have been obtained from the Tula Uplift in the western reach of the Eastern Kunlun Range (Figs. 2A and 9A; Robinson et al., 2003; Cheng et al., 2015a) and some Miocene to Quaternary volcanism occurred in the southwestern part of the range (Jolivet et al., 2003).

6.2 Provenance analyses of the Mesozoic-Cenozoic strata in the western Qaidam basin

6.2.1 Eboliang section

On a first order analysis, the detrital zircon U-Pb age spectrum obtained from the eight samples collected on the Eboliang section are largely similar (Fig. 7), suggesting that the provenance area was largely homogeneous through time. However, second-order variations in age cluster proportions among those samples record source changes during basin filling. In general, the detrital zircons U-Pb ages can be statistically subdivided into three major groups of Precambrian (spanning from ca. 2.8 Ga to 550 Ma), early to middle Paleozoic (peaks at 460~400 Ma) and late Paleozoic to Mesozoic (peaks at ca. 260~240 Ma) (Fig. 7). Aside from these major groups, the restricted early Neoproterozoic age group spanning from ca. 850 to ca. 1100 Ma with a peak age around 911 Ma (for example in sample E123, Lulehe Formation) represents the contribution from a distinctive source in the Altyn Tagh Range that will be discussed in details below.

Based on isopachs distribution, the Jurassic depocenter was situated farther south of the Eboliang section, which rules out any contribution from the Eastern Kunlun Range in the Jurassic sediments from that section (Fig. 9A; also see Fig. 3b in Meng et al., 2001). We thus consider that the paleocurrents in the northwestern Qaidam basin (Ritts and Biffi, 2000) probably provided sediments from mixed sources both in the Qilian Shan and Altyn Tagh Range. Based on $^{40}\text{Ar}/^{39}\text{Ar}$ thermochronology as well as zircon and apatite fission track analysis, previous

researches have reported Jurassic exhumation in the Altyn Tagh Range (Delville et al., 2001; Jolivet et al., 2001; Sobel et al., 2001; Wang et al., 2005). The detrital zircon age distribution of the Jurassic sediments in the Eboliang section (sample E122, Fig. 7D), is very similar to that of the Cenozoic samples, including early to middle Paleozoic zircons (peaks at 422 Ma; 56 % of total), late Paleozoic to Mesozoic zircons (peaks at ca. 253 Ma; 20 % of total) and a few Precambrian zircon (15 % of total). This similarity suggests that the source of the material deposited in the Eboliang region has been largely stable since the early Jurassic including both Permian granitoids and Ordovician to Devonian basement from either or both the Altyn Tagh Range and the Qilian Shan (Figs. 7A and 9).

The Paleogene depocenters of the Qaidam basin were consistently located south of the Eboliang section (see Fig. 12A in Yin et al., 2008b), again excluding material contribution from the Eastern Kunlun Range (Fig. 9A; Meng and Fang, 2008; Yin et al., 2008b; Mao et al., 2014). Paleocurrent directions obtained from conglomerate fabrics (clast imbrications) as well as heavy mineral assemblages suggest that the Paleogene clastic materials deposited in the northwestern Qaidam basin mainly derived from the Qilian Shan to the northeast and the Altyn Tagh Range to the north (Fig. 7; Wu et al., 2011; Fu et al., 2013; Jian et al., 2013). The age spectrum obtained from the three Paleogene samples (E123, EBLE, and E22) are characterized by early to middle Paleozoic zircons (peaks at ca. 414 to ca. 461 Ma; 20 % ~ 43 % of total), late Paleozoic to early Mesozoic zircons (peaks at ca. 246 Ma to ca. 255 Ma; 20 % ~ 23 % of total), and a progressively increasing proportion of Precambrian zircons

(from 23 % to 37 %).

The age spectrum of sample E123 displays a distinctive early Neoproterozoic age group (Fig. 7B; 1100 Ma to 850 Ma; peak at 911 Ma). The oscillatory zoning in CL images and the relatively high Th/U ratios (over 0.1, average value 0.3) of those early Neoproterozoic zircons in sample E123 indicate a clear magmatic origin (Fig. 10; Hanchar and Rudnick, 1995; Hoskin and Black, 2000; Corfu et al., 2003). As shown in Figure 9A, the early Neoproterozoic sources are few and small in the Altyn Tagh Range and Qilian Shan, but for a large magmatic complex north of the Anxi area. Wang et al. (2013) recently reported a mean zircon U-Pb crystallization age of ~910 Ma for these intrusions, consistent with the 911 Ma peak age in the Paleocene to early Eocene sample E123 (Fig. 9A). The enriched early Neoproterozoic zircon age groups (ca. 1100 to ca. 850 Ma) in sample E123 that accounts for over 20% of the total analysis represents a major difference from the age spectrum of the Jurassic and post-Paleocene samples. We suggest that this group indicates that the extensive early Neoproterozoic intrusion north of the Anxi area (Fig. 9A) formed a potential source for the Paleocene to early Eocene strata in the Eboliang section. According to the isopach map of the Paleocene to early Eocene Lulehe Formation provided by Yin et al. (2008b), the main depocenter was located south of the Eboliang section and the strata in the Eboliang section generally thicken southwards and westwards (Fig. 8). In addition, a sub-depocenter developed in the western Qaidam basin close to the early Neoproterozoic basement rocks in the Altyn Tagh Range (see Fig. 12A in Yin et al., 2008b). If the Eboliang section had been continuously situated in roughly the same

region as it occupies today (Fig. 2), the clastic materials derived from the early Neoproterozoic basement would thus not have been transported to the northwestern Qaidam basin. In addition, the few Neoproterozoic intrusions in the central and northern Qilian Shan region are not only far away from the Eboliang section, but also completely obstructed by the Paleocene uplift of the northern Qaidam thrust belt, clearly revealed by the seismic profiles in the southwestern flank of the Qilian Shan (Yin et al., 2008a). Based on these results combined with the generally south-directed paleoflows reported from the northern Qaidam basin (e.g. Fu et al., 2013; Jian et al., 2013) and the largely accepted hundreds of kilometers of Cenozoic offset along the lithospheric ATF (e.g. Wittlinger et al., 1998; Wang et al., 2006a; Yu et al., 2014; Cheng et al., 2015a, 2015b), we conclude that the Eboliang section must have been located much closer to the present day position of Anxi (Figs. 1 and 2). The single Paleogene zircon age observed in sample EBLE (Fig. 7C) is difficult to interpret since no Cenozoic magmatism has been reported in the Qaidam basin, Altyn Tagh Range or Qilian Shan. Our preferred interpretation is that this 57 Ma zircon was derived from the Cenozoic volcanic ash emitted from the volcanic centers exposed along the northern edge of the Qiangtang Block (e.g. Jolivet et al., 2003; Ding et al., 2007; Staisch et al., 2015).

The post-Oligocene depocenters are all located in the center of the Qaidam basin, south of the Eboliang section (Fig. 8; Meng and Fang, 2008; Yin et al., 2008b; Mao et al., 2014). This again excludes the possibility that the Neogene Eboliang sediments issued from the Eastern Kunlun Range to the south. Consistently, the generally

south-directed paleocurrents indicate that the sources of the post-Oligocene sediments in the Eboliang section were to the north, probably in the Altyn Tagh Range and the Qilian Shan (Wu et al., 2012b). However, the paleoflows changed from south-directed during the early Miocene, to generally southwest-directed during the Pliocene (Fig. 7). Unidirectional, south-directed paleocurrents prevail during the Quaternary (Fig. 7). Although Miocene to Quaternary strata are still dominated by early to middle Paleozoic zircons (peaks at ca. 444 Ma to ca. 401 Ma; 45 % ~ 48 % of total) and late Paleozoic to Mesozoic zircons (peaks at ca. 258 Ma to ca. 242 Ma; 39 % ~ 50 % of total), Precambrian zircons become scarce (3% ~ 9 % of total), especially in Pliocene and Quaternary strata (Fig. 7F, 7G and 7H). As shown in the 3D seismic profile (Fig. 11), growth strata that developed during the deposition of the early Miocene series most probably indicate intense tectonic deformation along the adjacent ATF. Based on sediment facies studies and provenance analysis of the Oligocene to Miocene strata in the Xorkol basin, Ritts et al. (2004) concluded that the Oligocene to early Miocene sequence coarsens upwards from low-energy playa and alluvial mudflat deposits to proximal alluvial fan deposits, again supporting early Miocene deformation within the Altyn Tagh Range. A similar transition has been observed in the Oligocene to Miocene strata along the northern flank of the Altyn Tagh Range (Yue et al., 2004a). In addition, Miocene deformation within the Altyn Tagh Range has been clearly registered by thermochronology and paleomagnetism data (e.g. Jolivet et al., 1999; Chen et al., 2001; Jolivet et al., 2001; Wang et al., 2006; Liu et al., 2007; Lu et al., 2014). We thus propose that the disappearance of the Precambrian zircon in the

post-Oligocene strata is linked to a major change in drainage pattern within the Altyn Tagh Range. The early Miocene deformation along the ATF led to the growth of a prominent topographic barrier, cutting off the link between the Precambrian source rocks in the Altyn Tagh Range and the Eboliang section area. Furthermore, the post-Oligocene samples (EBLN, EB1, ED1 and EE1) were collected in the southern part of the Eboliang fold structure that formed during or after the early Miocene (Figs. 2 and 11 A; e.g. Shang, 2001). Similarly to the material produced in the Altyn Tagh Range, the one sourced from the Qilian Shan was thus blocked east of the elevated growing structure and was not able to reach the deposition area to the south (Figs. 2, 9 and 11A). These results imply that the post-Oligocene clastic material deposited in the northwestern Qaidam basin was mainly derived from restricted local sources and/or recycling of the deformed pre- Oligocene strata.

6.2.2 Huatugou section

Along this section the detrital zircons U-Pb ages are again statistically subdivided into three major populations of Precambrian (spanning from ca. 3.2 Ga to 600 Ma), early to middle Paleozoic (peaks at 460~400 Ma) and late Paleozoic to Mesozoic (peaks at ca. 290~210 Ma) (Fig. 8). The Jurassic to Pleistocene basin depocenters were consistently situated farther southeast of the Huatugou section, which rules out any material contribution from the Qilian Shan to the Huatugou section (Fig. 9A; Meng and Fang, 2008; Yin et al., 2008b; Mao et al., 2014).

Late Paleozoic - early Mesozoic granitic intrusions are widespread in the Altyn Tagh and Eastern Kunlun ranges, however, most of the early Mesozoic intrusions

(younger than ca. 260 Ma) are confined to the Eastern Kunlun Range just south of our sample sites (Fig. 9A). This distinctive pluton distribution, together with the early Neoproterozoic zircons (1100~850 Ma) issued from the remarkably large intrusion in the Altyn Tagh Range (Fig. 9A; Wang et al., 2013; Cheng et al., 2015a) can be used as provenance signatures to identify the source of detrital zircons within our nine sandstone samples.

The poorly sorted, angular grains in the Jurassic samples suggest deposition from a proximal source (see Figs. 5A and 5B in Cheng et al., 2015a). The zircon ages spectrum of the Jurassic sample (CSL3, Fig. 8A) is characterized by an exceptional bimodal distribution (Permian-Triassic and Paleozoic groups). The zircons with ages younger than 260 Ma account for 15.5 % of the total number of analysis, suggesting that the Eastern Kunlun Range was a significant source of material for the Jurassic strata in the Huatugou section. This result is consistent with the general north- to west- directed paleoflows (Fig. 8A). The few Precambrian zircons (no early Neoproterozoic zircons) as well as the few south-directed paleocurrents might indicate limited contribution from the Altyn Tagh Range to the north, which suggests a limited topographic relief within that range, locally exposing basement rocks (Delville et al., 2001; Sobel et al., 2001; Ritts and Biffi, 2000).

In the Paleocene-early Eocene sample (S23) and the middle Eocene sample (HTG-E), the age spectrums are still largely bimodal, with about 15% of < 260 Ma zircons, suggesting that the Eastern Kunlun Range was still a significant source for these samples. This assumption is consistent with the N-NE-directed paleoflows

601 observed in the Paleocene to middle Eocene series (Fig. 8B and 8C; Meng and Fang,
602 2008; Fu et al., 2013; Li et al., 2015). On the other hand, the increasing proportion of
603 Precambrian zircons (over 3% of early Neoproterozoic zircons in both samples)
604 indicates that the Altyn Tagh Range was gradually becoming a significant source of
605 clastic material for the Paleocene to middle Eocene strata in the Huatugou section.

606 The late Eocene to Oligocene sample (QX1-2) displays a significant change in the
607 detrital zircon age distribution pattern (Fig. 8C, 8D and 8E): the major Mesozoic age
608 peak observed in the previous samples becomes nearly negligible with the proportion
609 of < 260 Ma zircons decreasing dramatically (from 13.6% in sample S23 and 18.4%
610 in sample HTG-E to 3.1% in sample QX1-2). In addition, the paleocurrent directions
611 changed abruptly from generally north-directed in the Paleocene to middle Eocene
612 strata to south-directed in the late Eocene strata (Fig. 8C and 8D). This implies a
613 major shift in the source region from the Eastern Kunlun Range to the Altyn Tagh
614 Range during the middle to late Eocene time. The proportion of early Neoproterozoic
615 zircons gradually increased (e.g. from 3.1% in sample HTG-E to 5.1% in sample
616 QX1-2) further attesting that basement rocks in the Altyn Tagh Range served as a
617 significant source for the sediments in the Huatugou section.

618 Recent sedimentology and paleomagnetic studies demonstrated that the relatively
619 rigid Qaidam basin, together with the western segment of the Eastern Kunlun Range,
620 have been transported northeastward along the ATF during the Cenozoic (Wang et al.,
621 2006a; Yu et al., 2014; Cheng et al., 2015a, 2015b). This displacement occurred
622 without obvious basin-scale vertical axis rotation with respect to the Eurasia Plate

since the early Eocene (Dupont-Nivet et al., 2002; Yu et al., 2014). We suggest that this sharp change in the detrital zircon age distribution pattern was directly related to the onset of large-scale left-lateral strike-slip displacement along the ATF. Except for the still slightly increasing proportion of Precambrian ages, the detrital zircon U-Pb age distribution registered in the Oligocene sample (QX1-1) is similar to that in the late Eocene sample (QX1-2), suggesting a relatively stable drainage pattern from the late Eocene to the Oligocene (Fig. 8D and 8E).

The age spectrum of the early Miocene sample (HTG-N) is again characterized by a major Paleozoic age peak and a minor Mesozoic age sub-peak (Fig. 8F). However, the amount of Precambrian zircons decreased significantly (early Neoproterozoic ages only represent 1.1% of the total analysis). A similar decrease in the proportion of Precambrian zircons was also registered in the early Miocene sample of the Eboliang section (sample EBLN in Fig. 7E), indicating a regional change in the source area and/or drainage pattern throughout the Altyn Tagh Range. As already mentioned in the provenance analysis of samples from the Eboliang section, we suggest that the regional early Miocene deformation within the Altyn Tagh Range cut off the link between the Precambrian source rocks in the Altyn Tagh and Qilian Shan ranges and the depositional area in the Huatugou and Eboliang sections. The early Miocene clastic material in these two sections was mainly derived from more restricted local source regions and recycling of the deformed Paleocene to Oligocene strata.

Detrital zircon age distributions in the middle Miocene to Pleistocene samples

(CSL4, SZG1 and CSL5) are similar, characterized by a major middle Paleozoic age peak, a minor but consistent Mesozoic age peak as well as few Precambrian zircons (Fig. 8G, 8H and 8I). Based on seismic profile interpretation, the fold structures developing perpendicular to the Altyn Tagh fault trend (e.g. the Yingxiongling structure, Fig. 2) initiated during or after the Oligocene (Yin et al., 2007; Yin et al., 2008a; Wu et al., 2014). After the Oligocene onset of deformation, the Huatugou section, located in the northernmost part of the prominent Yingxiongling uplift structure, was gradually isolated from the main Qaidam basin. A complex, local drainage pattern developed while recycling of the deformed Paleocene to Oligocene strata served as a major material source (Fig. 8).

6.3 Tectonics implications

6.3.1 Source to sink relation between the Qaidam basin and the Altyn Tagh Range

Provenance analysis results obtained from the two studied sections along the ATF reveal that the source to sink relation between the Qaidam basin and the Altyn Tagh Range underwent a three stages evolution:

During the Paleocene and early Eocene, the Eboliang and Huatugou sections were respectively located near the present day position of Anxi and Tula (Figs. 2A and 12A). The basement rocks in the Altyn Tagh Range (including the early Neoproterozoic intrusion north of the Anxi area) served as the major source area for the Eboliang section while a part of the material deposited in the Huatugou section derived from the Eastern Kunlun Range.

By early to middle Eocene, left-lateral strike-slip movement along the ATF

initiated, resulting in gradually northeastward migration of the Qaidam basin (including the Eboliang and Huatugou sections). This migration ultimately disconnected the Eboliang section from the early Neoproterozoic source north of the Anxi area (Fig. 12B), as attested by the decreasing proportion of early Neoproterozoic zircons (from 19.4% in Paleocene to early Eocene sample E123, to 5.3% in middle-late Eocene sample EBLE and 3.6% in the Oligocene sample E22). Similarly, the Huatugou section gradually moved away from the Tula area towards Anxi (Fig. 12B). This led to a decreasing proportion of < 260 Ma zircons (from ca. 15% in the Paleocene to middle Eocene samples S23 and HTG-E to ca. 3% in the late Eocene sample QX1-2; Fig. 8B, 8C and 8D) and to an abrupt change in paleocurrent directions (Fig. 8C and 8D). In parallel, the gradual increase in early Neoproterozoic zircons (from ca. 3% in middle Eocene sample HTG-E to 6.5% in Oligocene sample QX1-1; Fig. 8C, 8D and 8E) suggests that the Huatugou section might be close to the early Neoproterozoic intrusives north of the Anxi area during the middle Eocene to Oligocene.

On seismic profiles BB' and CC' (trending perpendicular to the Altyn Tagh Range; Figs. 2, 11A and 11B), several basement-involved thrust faults offset the Mesozoic strata and die out in the Eocene and Oligocene strata. In addition, the Eocene to Oligocene strata slightly thicken towards the depocenter exhibiting a growth-strata structure (Fig. 11B). The oldest unit affected by growth strata is the middle Eocene lower Xiaganchaigou Fm., suggesting that the deformation along the ATF initiated around that time. Paleocurrents measurements and provenance analysis of

conglomerates in the Xorkol basin (Fig. 2) and western Qaidam basin indicate that uplift along the central segment of the ATF occurred prior to the Oligocene (Yue et al., 2001; Ritts et al., 2004; Wu et al., 2012a, 2012b), probably during the early to middle Eocene.

Since the early Miocene, the Eboliang and Huatugou sections continuously migrated northeastwards along the left-lateral strike-slip ATF. However, the scarcity of the Precambrian zircons, especially the early Neoproterozoic zircons, in the post-Oligocene sediments (e.g. sample EBLN in the Eboliang section and sample HTG-N in the Huatugou section) indicates a major change in the drainage pattern leading to disruption of the source to sink relation between the Altyn Tagh Range and the western Qaidam basin. On seismic profiles BB' and CC' (Fig. 11A and 11B), generally northward-tapering growth strata within the Miocene to Pliocene deposits are well-developed, likely associated with the left-lateral faulting along the ATF and deformation within the Altyn Tagh Range since the early Miocene. The post-Oligocene deformation and uplift within the Altyn Tagh Range and northwestern Qaidam basin largely modified the regional drainage pattern. Most of the material derived from the erosion of the Altyn Tagh Range was directed towards the Tarim basin to the north (Fig. 12C; Yue et al., 2004b). The post-Oligocene clastic materials deposited in the western Qaidam basin was derived from local sources including recycling of the deformed Paleocene to Oligocene strata.

6.3.2 Implications for the Miocene deformation within the Altyn Tagh Range

In this study, the disappearance of an early Neoproterozoic component in the

711 detrital zircon age spectra of the Miocene strata both in the Eboliang and Huatugou
712 sections as well as the Miocene initiation of growth strata within the northwestern
713 Qaidam basin are indicative of enhanced tectonic activity within the Altyn Tagh
714 Range and Qaidam basin since the Miocene. This finding is consistent with the results
715 of several previous multi-faceted studies including: thermochronology on the
716 basement rocks within the Altyn Tagh Range (e.g. Chen et al., 2001; Jolivet et al.,
717 1999, 2001; Sobel et al., 2001; Zhang et al., 2012); paleomagnetism studies on the
718 Cenozoic strata within the northwestern Qaidam basin (e.g. Zhang et al., 2013b; Lu et
719 al., 2014; Chang et al., 2015); basin-scale seismic profile interpretation (Meng and
720 Fang, 2008; Yin et al., 2008a, 2008b; Wang et al., 2010; Cheng et al., 2015a);
721 provenance analyses along the Altyn Tagh Range (e.g. Yue et al., 2001; Yin et al.,
722 2002; Wu et al., 2012a, 2012b) and synthetized tectonic analysis (e.g. Burchfiel et
723 al., 1989; Meyer et al., 1998; Fu et al., 2015). In addition, the Miocene deformation
724 phase has been widely recognized in the northern and eastern Tibetan plateau,
725 including the Eastern Kunlun Range, the Altyn Tagh Range, the Longmen Shan and
726 the Qilian Shan (e.g., Kirby et al., 2002; Clark et al., 2004; Ding et al., 2004; Ritts et
727 al., 2004; Sun et al., 2005; Duvall et al., 2013; Yuan et al., 2013; Cheng et al., 2014,
728 2015a, 2016; Lease, 2014; Chang et al., 2015; Jolivet et al., 2015), which may reflect
729 a critical period in the growth process of the Tibetan plateau.

730 6.3.3 Implications for the Cenozoic offset of the ATF

731 The Eboliang and Huatugou sections record the northward migration of the
732 Qaidam basin and can be used as two piercing points for defining the Cenozoic offset

of the ATF. In addition, the early Neoproterozoic intrusion in the Altyn Tagh Range strikes NEE-SWW along the Altyn Tagh fault, and has an elongation of ca. 80 km. The Eboliang section is located south of this 80 km wide early Neoproterozoic intrusion and separated from the intrusion by the ATF. Considering the actual 360 km gap between the center of this early Neoproterozoic intrusion and the Eboliang section, an average of 360 ± 40 km offset on the Altyn Tagh fault can be estimated. A similar ~360 km offset is obtained by correlating the Huatugou section to the Mesozoic plutons south of the Tula area. We concede that, from analyzing detrital zircon spectra of Mesozoic to Cenozoic strata, the Eboliang-Anxi piercing point is much more convincing than the Tula-Huatugou piercing point. However, by correlating the lithology and sedimentary features of the Cenozoic strata and analyzing the detrital zircon U-Pb ages from Mesozoic strata in both the Tula and Huatugou sections, we previously found potential links between both sections (Cheng et al., 2015a). We thus considered that the Tula section used to be a part of the Qaidam basin and was left behind due to the northeastward migration of the Qaidam basin. Two faults which have been indentified by previous research (Figs. 1, 2 and 12; e.g. Cowgill, et al., 2003) are acting as boundaries between the detached Qaidam basin and the relic Tula sub-basin (Cheng et al., 2015a). Since the early Eocene, faulting on the ATF induced these NE-SW–trending branch faults, now covered by Quaternary deposits. Northwards migration of the rigid Qaidam basin left the Tula section and parts of the Kunlun basement behind (Cheng et al., 2015a). Moreover, considering the strong mechanical behaviour of the rigid Qaidam basin, the Huatugou section should be

approximately located near the present-day position of the Tula section if we restore the horizontal offset along the AFT based on the convincing Eboliang-Anxi piercing point. Consequently, we suggest a 360 ± 40 km offset along the ATF, which contradicts the extraordinary ~ 1200 km offset estimation (CSBS, 1992), but is in excellent agreement with independent, albeit indirect piercing points suggesting 300 to 400 km of displacement along the Altyn Tagh Range: (1) 400 ± 60 km offset based on Jurassic lacustrine shorelines along the northern and southern sides of the Altyn Tagh fault (Ritts and Biffi, 2000), (2) ~ 375 km offset of the Cambrian magmatic arc in the Qilian Shan and the Altyn Tagh Range (Gehrels et al., 2003a), (3) $\sim 375 \pm 25$ km offset of pre-Oligocene strata between the Qaidam and Xorkol basins (Yue et al., 2001), (4) $\sim 360 \pm 40$ km offset estimation based on source to sink relation between the Xorkol basin and the North Qilian Shan (Yue et al., 2005), (5) 350~400 km offset based on eclogite, ophiolite and blueschist facies units in the Altyn Tagh Range and Qilian Shan (Yang et al., 2001; Zhang et al., 2001), (6) ~ 350 km offset based on similar early to middle Jurassic cooling zones on both side of the Altyn Tagh fault (Sobel et al., 2001). This large-scale lithospheric offset along the ATF would be totally accommodated by crustal deformation within the terrane to the east, especially by the NE-SW shortening and eastward extrusion along the faults within the Qilian Shan (e.g. Yin et al., 2002, 2008a; Cheng et al., 2015b). The confirmation of the hundreds of kilometers of displacement along the strike-slip ATF again supports the eastward extrusion of the Tibetan plateau driven by the India-Asia collision.

7. Conclusion

Detrital zircon U-Pb age patterns evolution from two Jurassic to Pleistocene stratigraphic sections in the northwestern Qaidam basin associated to high-quality seismic-reflection profiles revealed that:

1. During the Paleocene to early Eocene, the Eboliang and Huatugou sections were respectively located near the present-day position of the Anxi and Tula areas. The basement rocks in the Altyn Tagh Range served as the significant source for the sediments to the Eboliang section, while the material deposited in the Huatugou section mostly derived from the Eastern Kunlun Range.
2. Left-lateral strike-slip movement along the ATF initiated during the early-middle Eocene, resulting in gradual northeastward migration of the Qaidam basin.
3. The post-Oligocene deformation within the Altyn Tagh Range and northwestern Qaidam basin strongly modified the regional drainage pattern. By Oligocene times, most of the material issued from the erosion of the Altyn Tagh Range was directed towards the Tarim basin to the north. The post-Oligocene clastic materials in the western Qaidam basin was mainly derived from local sources largely including recycling of the deformed Paleocene to Oligocene strata.
4. Using the Eboliang and Huatugou sections as piercing points, we estimate a 360 ± 40 km offset along the ATF.

Acknowledgements

This work was supported by the Oil and Gas Scientific and Technologic Major Programs of China (Grant ZX05003) and the China Scholarship Council – Campus France Egide Cai Yuanpei program (201404490136; N° 32393WL (2014)). We gratefully acknowledge the assistance of Shuwei Guan, Qinyang Meng, Ruiying Chen, Chuanmin Zhou, Suping Zhou, Xiangjiang Yu, Xiang Cheng, Wei Du, Runchao Liu and Tuo Zhang during field work. We thank Li Su of the Geological Lab Center, China University of Geosciences (Beijing), for her help during detrital zircon analysis and to Chaodong Wu, Chuanwu Wang, Yan Chen for their helpful suggestions on the Cenozoic evolution of the Qaidam basin. Communications with Michel Balleve, Sylvie Bourquin, Hervé Régnault, Baofu Han and Zhicheng Zhang have sharpened the discussion on the appropriate piercing points. We particularly thank Science Editor Jean-Philippe Avouac and two anonymous reviewers for their careful and insightful comments. We gratefully acknowledge the journal manager, Iswarya Samikannu, for handling our manuscript.

Support Information

Appendix A. U-Pb analysis of detrital zircons from the 11 sandstone samples.

Appendix B. Detailed description of the U–Pb geochronology results.

References

- Andersen, T., 2002. Correction of common lead in U-Pb analyses that do not report ^{204}Pb . *Chemical Geology* 192, 59-79.
- Arnaud, N., Tapponnier, P., Roger, F., Brunel, M., Scharer, U., Chen, W., Xu, Z., 2003. Evidence for Mesozoic shear along the western Kunlun and Altyn-Tagh fault, northern Tibet (China). *Journal of Geophysical Research: Solid Earth* (1978-2012) 108.
- Avouac, J.P., Tapponnier, P., 1993. Kinematic model of active deformation in central Asia. *Geophysical Research Letters* 20, 895-898.
- Black, L.P., Kamo, S.L., Allen, C.M., Aleinikoff, J.N., Davis, D.W., Korsch, R.J., Foudoulis, C., 2003. TEMORA 1: a new zircon standard for Phanerozoic U-Pb geochronology. *Chemical Geology* 200, 155-170.
- Bovet, P.M., Ritts, B.D., Gehrels, G., Abbink, A.O., Darby, B., Hourigan, J., 2009. Evidence of Miocene crustal shortening in the north Qilian Shan from Cenozoic stratigraphy of the western Hexi Corridor, Gansu Province, China. *American Journal of Science* 309, 290-329.
- Braitenberg, C., Wang, Y., Fang, J., Hsu, H., 2003. Spatial variations of flexure parameters over the Tibet–Qinghai plateau. *Earth and Planetary Science Letters* 205, 211-224.
- Burchfiel, B.C., Quidong, D., Molnar, P., Royden, L., Yipeng, W., Peizhen, Z., Weiqi, Z., 1989. Intracrustal detachment within zones of continental deformation. *Geology* 17, 748-752.

840 Chang, H., Li, L., Qiang, X., Garzione, C.N., Pullen, A., An, Z., 2015.
841 Magnetostratigraphy of Cenozoic deposits in the western Qaidam Basin and its
842 implication for the surface uplift of the northeastern margin of the Tibetan Plateau.
843 Earth and Planetary Science Letters 430, 271-283.

844 Chen, Z.L., Zhang, Y.Q., Wang, X.F., Chen, X.H., 2001. Fission Track Dating of
845 Apatite Constrains on the Cenozoic Uplift of the Altyn Tagh Mountain. Acta
846 Geoscientia Sinica 22, 413-418 (in Chinese with English abstract).

847 Chen, X.H., Wang, X.F., E. George, G., Yang, Y., Qin, H., Chen, Z.L., Yang, F., Chen,
848 B.L., Li, X.Z., 2004. Early Paleozoic Magmatism and Gold Mineralization in the
849 Northern Altun, NW China. Acta Geologica Sinica-English Edition 78, 515-523.

850 Chen, X., Gehrels, G., Yin, A., Zhou, Q., Huang, P., 2015. Geochemical and
851 Nd–Sr–Pb–O isotopic constrains on Permo–Triassic magmatism in eastern Qaidam
852 Basin, northern Qinghai-Tibetan plateau: Implications for the evolution of the
853 Paleo-Tethys. Journal of Asian Earth Sciences 114, Part 4, 674-692.

854 Chen, Y., Gilder, S., Halim, N., Cogné, J.P., Courtillot, V., 2002. New paleomagnetic
855 constraints on central Asian kinematics: Displacement along the Altyn Tagh fault and
856 rotation of the Qaidam Basin. Tectonics 21, 1042.

857 Cheng, F., Jolivet, M., Fu, S., Zhang, Q., Guan, S., Yu, X., Guo, Z., 2014. Northward
858 growth of the Qimen Tagh Range: A new model accounting for the Late Neogene
859 strike-slip deformation of the SW Qaidam Basin. Tectonophysics 632, 32-47.

860 Cheng, F., Guo, Z., Jenkins, H.S., Fu, S., Cheng, X., 2015a. Initial rupture and
861 displacement on the Altyn Tagh fault, northern Tibetan Plateau: Constraints based on

862 residual Mesozoic to Cenozoic strata in the western Qaidam Basin. *Geosphere* 11,
863 921-942.

864 Cheng, F., Jolivet, M., Dupont-Nivet, G., Wang, L., Yu, X., Guo, Z., 2015b. Lateral
865 extrusion along the Altyn Tagh Fault, Qilian Shan (NE Tibet): insight from a 3D
866 crustal budget. *Terra Nova* 27, 416-425.

867 Cheng, F., Fu, S., Jolivet, M., Zhang, C., Guo, Z., 2016. Source to sink relation
868 between the Eastern Kunlun Range and the Qaidam Basin, northern Tibetan Plateau,
869 during the Cenozoic. *Geological Society of America Bulletin* 128, 258-283.

870 Clark, M., Schoenbohm, L., Royden, L., Whipple, K., Burchfiel, B., Zhang, X., Tang,
871 W., Wang, E., Chen, L., 2004. Surface uplift, tectonics, and erosion of eastern Tibet
872 from large-scale drainage patterns. *Tectonics* 23, TC1006.

873 Corfu, F., Hanchar, J.M., Hoskin, P.W., Kinny, P., 2003. Atlas of zircon textures.
874 *Reviews in mineralogy and geochemistry* 53, 469-500.

875 Cowgill, E., Yin, A., Harrison, T.M., Xiao-Feng, W., 2003. Reconstruction of the
876 Altyn Tagh fault based on U-Pb geochronology: Role of back thrusts, mantle sutures,
877 and heterogeneous crustal strength in forming the Tibetan Plateau. *Journal of*
878 *Geophysical Research* 108, 2346.

879 CSBS (Chinese State Bureau of Seismology), 1992. The Altyn Tagh Active Fault
880 System. Seismology Publishing House, Beijing, pp. 1-319 (in Chinese).

881 Dai, J., Wang, C., Hourigan, J., Santosh, M., 2013. Multi-stage tectono-magmatic
882 events of the Eastern Kunlun Range, northern Tibet: Insights from U-Pb
883 geochronology and (U-Th)/He thermochronology. *Tectonophysics* 599, 97-106.

884 Darby, B.J., Ritts, B.D., Yue, Y., Meng, Q., 2005. Did the Altyn Tagh fault extend
885 beyond the Tibetan Plateau? *Earth and Planetary Science Letters* 240, 425-435.

886 Delville, N., Arnaud, N., Montel, J.-M., Roger, F., Brunel, M., Tapponnier, P., Sobel,
887 E., 2001. Paleozoic to Cenozoic deformation along the Altyn Tagh fault in the Altun
888 Shan massif area, eastern Qilian Shan, northeastern Tibet, China. *Geological Society*
889 *of America Memoirs* 194, 269-292.

890 Deng, T., Wang, X.M., 2004a. New material of the Neogene rhinocerotids from the
891 Qaidam Basin in Qinghai, China. *Vert. PalAsiat.* 42, 216–229.

892 Deng, T., Wang, X.M., 2004b. Late Miocene Hipparion (Equidae, Mammalia) of
893 eastern Qaidam Basin in Qinghai, China. *Vert. PalAsiat.* 42, 316–333.

894 Ding, G., Chen, J., Tian, Q., Shen, X., Xing, C., Wei, K., 2004. Active faults and
895 magnitudes of left-lateral displacement along the northern margin of the Tibetan
896 Plateau. *Tectonophysics* 380, 243-260.

897 Ding, L., Kapp, P., Yue, Y., Lai, Q., 2007. Postcollisional calc-alkaline lavas and
898 xenoliths from the southern Qiangtang terrane, central Tibet. *Earth and Planetary*
899 *Science Letters* 254, 28-38.

900 Dong, Z., Pei, X., Gu, P., Zhang, H., Chen, R., Zhang, X., 2014. Geochronology,
901 Geochemistry, and Hf Isopleth of Yanchangbeishan Granodiorite of Lenghu Area in
902 Qinghai. *Nonwestern Geology* 47 (4), 141-155 (in Chinese with English abstract).

903 Dong, Z., Gu, P., Chen, R., Zhang, X., Zhang, H., 2015. Geochronology,
904 Geochemistry, and Hf Isopleth of Yanchangbeishan Adamellite of Lenghu Area in
905 Qinghai. *Earth Science-Journal of China University of Geosciences* 40, 130-144 (in

Chinese with English abstract).

Duvall, A.R., Clark, M.K., Kirby, E., Farley, K.A., Craddock, W.H., Li, C., Yuan, D.-Y., 2013. Low-temperature thermochronometry along the Kunlun and Haiyuan Faults, NE Tibetan Plateau: Evidence for kinematic change during late-stage orogenesis. *Tectonics* 32, 1190-1211.

Dupont-Nivet, G., Butler, R.F., Yin, A., Chen, X., 2002. Paleomagnetism indicates no Neogene rotation of the Qaidam Basin in northern Tibet during Indo-Asian collision. *Geology* 30, 263.

Dupont-Nivet, G., Lippert, P.C., Van Hinsbergen, D.J.J., Meijers, M.J.M., Kapp, P., 2010. Palaeolatitude and age of the Indo–Asia collision: palaeomagnetic constraints. *Geophysical Journal International* 182, 1189-1198.

England, P., Houseman, G., 1989. Extension during continental convergence, with application to the Tibetan Plateau. *Journal of Geophysical Research: Solid Earth* (1978–2012) 94, 17561-17579.

Fang, X., Zhang, W., Meng, Q., Gao, J., Wang, X., King, J., Song, C., Dai, S., Miao, Y., 2007. High-resolution magnetostratigraphy of the Neogene Huaitoutala section in the eastern Qaidam Basin on the NE Tibetan Plateau, Qinghai Province, China and its implication on tectonic uplift of the NE Tibetan Plateau. *Earth and Planetary Science Letters* 258, 293-306.

Fedo, C.M., Sircombe, K.N., Rainbird, R.H., 2003. Detrital zircon analysis of the sedimentary record. *Reviews in mineralogy and geochemistry* 53, 277-303.

Fu, L., Guan, P., Zhao, W. Y., Wang, M., Zhang, Y., Lu, J. W., 2013. Heavy mineral

928 feature and provenance analysis of Paleogene Lulehe Formation in Qaidam Basin.
 929 *Acta Petrologica Sinica* 29, 2867-2875 (in Chinese with English abstract).

930 Fu, S., Ma, D., Guo, Z., Cheng, F., 2015. Strike-slip superimposed Qaidam Basin and
 931 its control on oil and gas accumulation, NW China, *Petroleum Exploration and*
 932 *Development* 42, 779-789.

933 Gao, J., Li, S., Dai, S., Li, A., Peng, Y., 2009. Constraints of tectonic evolution in
 934 provenance from detrital zircon fission-track data of Cenozoic strata of Xichagou
 935 district in western Qaidam basin. *Journal of Lanzhou University (Natural Sciences)* 45,
 936 1-7 (in Chinese with English abstract).

937 Gehrels, G.E., Yin, A., Wang, X.F., 2003a. Magmatic history of the northeastern
 938 Tibetan Plateau. *Journal of Geophysical Research* 108, 2423.

939 Gehrels, G.E., Yin, A., Wang, X.F., 2003b. Detrital-zircon geochronology of the
 940 northeastern Tibetan plateau. *Geological Society of America Bulletin* 115, 881-896.

941 Gehrels, G., Kapp, P., DeCelles, P., Pullen, A., Blakey, R., Weislogel, A., Ding, L.,
 942 Guynn, J., Martin, A., McQuarrie, N., 2011. Detrital zircon geochronology of
 943 pre-Tertiary strata in the Tibetan-Himalayan orogen. *Tectonics* 30, TC5016.

944 Gehrels, G., 2014. Detrital Zircon U-Pb Geochronology Applied to Tectonics. *Annual*
 945 *Review of Earth and Planetary Sciences* 42, 127-149.

946 Guo, J.J., Li, H.K., 1999. Angular Unconformity between the Huashishan Group and
 947 Huangzhong Group in the Eastern Mid-Qilian Massif: Identification and Implications.
 948 *Progress in Precambrian Research* 22, 47-52.

949 Hanchar, J., Rudnick, R., 1995. Revealing hidden structures: the application of

950 cathodoluminescence and back-scattered electron imaging to dating zircons from
951 lower crustal xenoliths. *Lithos* 36, 289-303.

952 Hoskin, P., Black, L., 2000. Metamorphic zircon formation by solid-state
953 recrystallization of protolith igneous zircon. *Journal of metamorphic Geology* 18,
954 423-439.

955 Heermance, R.V., Pullen, A., Kapp, P., Garzione, C.N., Bogue, S., Ding, L., Song, P.,
956 2013. Climatic and tectonic controls on sedimentation and erosion during the
957 Pliocene–Quaternary in the Qaidam Basin (China). *Geological Society of America*
958 *Bulletin* 125, 833-856.

959 Huang, H., Huang, Q., Ma, Y., 1996. *Geology of Qaidam Basin and its Petroleum*
960 *Prediction*. Geological Publishing House, Beijing, pp. 1–257 (in Chinese with English
961 abstract).

962 HGSI (Hunan Geological Survey Institute), 2003. *The Report of Regional Geological*
963 *Survey of Qiemo at Scale 1:250,000*. China Industry Press, Beijing, pp. 1-281 in
964 Chinese with English abstract).

965 Huo, G.M., 1990. *Petroleum Geology of China: Oil Fields in Qianghai and Xizang*, 14.
966 Chinese Petroleum Industry Press, Beijing, pp. 1-483 (in Chinese with English
967 abstract).

968 IGSQP (Institute of Geological Survey of Qinhai Province), 2004. *The Report of*
969 *Regional Geological Survey at Scale 1:250000*. P.R.C. China Industry Press, Beijing,
970 pp. 1–386 (in Chinese).

971 Jian, X., Guan, P., Zhang, D.W., Zhang, W., Feng, F., Liu, R.J., Lin, S.D., 2013.

972 Provenance of Tertiary sandstone in the northern Qaidam basin, northeastern Tibetan
973 Plateau: Integration of framework petrography, heavy mineral analysis and mineral
974 chemistry. *Sedimentary Geology* 290, 109-125.

975 Jolivet, M., Roger, F., Arnaud, N., Brunel, M., Tapponnier, P., Seward, D., 1999.
976 Histoire de l'exhumation de l'Altun Shan: indications sur l'âge de la subduction du
977 bloc du Tarim sous le système de l'Altyn Tagh (Nord Tibet). *Comptes Rendus de*
978 *l'Académie des Sciences-Series IIA-Earth and Planetary Science* 329, 749-755.

979 Jolivet, M., Brunel, M., Seward, D., Xu, Z., Yang, J., Roger, F., Tapponnier, P.,
980 Malavieille, J., Arnaud, N., Wu, C., 2001. Mesozoic and Cenozoic tectonics of the
981 northern edge of the Tibetan plateau: fission-track constraints. *Tectonophysics* 343,
982 111-134.

983 Jolivet, M., Brunel, M., Seward, D., Xu, Z., Yang, J., Malavieille, J., Roger, F.,
984 Leyreloup, A., Arnaud, N., Wu, C., 2003. Neogene extension and volcanism in the
985 Kunlun Fault Zone, northern Tibet: New constraints on the age of the Kunlun Fault.
986 *Tectonics* 22, 1052.

987 Jolivet, M., Roger, F., Xu, Z.Q., Paquette, J.L., Cao, H., 2015. Mesozoic–Cenozoic
988 evolution of the Danba dome (Songpan Garzê, East Tibet) as inferred from
989 LA-ICPMS U–Pb and fission-track data. *Journal of Asian Earth Sciences* 102,
990 180-204.

991 Ke, X., Ji, J., Zhang, K., Kou, X., Song, B., Wang, C., 2013. Magnetostratigraphy and
992 Anisotropy of Magnetic Susceptibility of the Lulehe Formation in the Northeastern
993 Qaidam Basin. *Acta Geologica Sinica-English Edition* 87, 576-587.

994 Kirby, E., Reiners, P.W., Krol, M.A., Whipple, K.X., Hodges, K.V., Farley, K.A., Tang,
 995 W.Q., Chen, Z.L., 2002. Late Cenozoic evolution of the eastern margin of the Tibetan
 996 Plateau: Inferences from $^{40}\text{Ar}/^{39}\text{Ar}$ and (U-Th)/He thermochronology. *Tectonics* 21,
 997 1001.
 998 Lease, R.O., 2014. Cenozoic mountain building on the northeastern Tibetan Plateau.
 999 Geological Society of America Special Papers 507, 115-127.
 1000 Li, H.B., Yang, J.S., Xu, Z.Q., Sun, Z.M., Tapponnier, P., van der Woerd, J., Meriaux,
 1001 A.-S., 2006. The constraint of the Altyn Tagh fault system to the growth and rise of
 1002 the northern Tibetan Plateau. *Earth Science Frontiers* 13, 59-79 (in Chinese with
 1003 English abstract).
 1004 Li, W., Neubauer, F., Liu, Y.J., Genser, J., Ren, S.M., Han, G.Q., Liang, C.Y., 2013.
 1005 Paleozoic evolution of the Qimantagh magmatic arcs, Eastern Kunlun Mountains:
 1006 constraints from zircon dating of granitoids and modern river sands. *Journal of Asian*
 1007 *Earth Sciences* 77, 183-202.
 1008 Li, L., Guo, Z., Guan, S., Zhou, S., Wang, M., Fang, Y., Zhang, C., 2015. Heavy
 1009 mineral assemblage characteristics and the Cenozoic paleogeographic evolution in
 1010 southwestern Qaidam Basin. *Science China Earth Sciences* 58, 859-875.
 1011 Liu, Y. H., Mo, X. X., Yu, X. H., Zhang, X. T. and Xu, G. W., 2006. Zircon SHRIMP
 1012 U-Pb dating of the Jingren granite, Yemaquan region of the east Kunlun and its
 1013 geological significance: *Acta Petrologica Sinica* 22, 2457-2463 (in Chinese with
 1014 English abstract).
 1015 Liu, Y.J., Neubauer, F., Genser, J., Ge, X.H., Takasu, A., Yuan, S.H., Chang, L.H., Li,

1016 W.M., 2007. Geochronology of the initiation and displacement of the Altyn
 1017 Strike-Slip Fault, western China. *Journal of Asian Earth Sciences* 29, 243-252.

1018 Liu, D., Jolivet, M., Yang, W., Zhang, Z., Cheng, F., Zhu, B., Guo, Z., 2013. Latest
 1019 Paleozoic–Early Mesozoic basin–range interactions in South Tian Shan (northwest
 1020 China) and their tectonic significance: Constraints from detrital zircon U–Pb ages.
 1021 *Tectonophysics* 599, 197-213.

1022 Liu, D., Cheng, F., Guo, Z., Jolivet, M., Song, Y., 2015. Lahar facies of the Latest
 1023 Paleozoic Arbasay Formation: Geomorphological characters and paleoenvironment
 1024 reconstruction of Northern Tian Shan, NW China. *Journal of Asian Earth Sciences*
 1025 113, Part 1, 282-292.

1026 Long, X., Yuan, C., Sun, M., Kröner, A., Zhao, G., 2014. New geochemical and
 1027 combined zircon U–Pb and Lu–Hf isotopic data of orthogneisses in the northern Altyn
 1028 Tagh, northern margin of the Tibetan plateau: Implication for Archean evolution of the
 1029 Dunhuang Block and crust formation in NW China. *Lithos* 200–201, 418-431.

1030 Lu, S.N., 2002. Preliminary Study of Precambrian Geology in the North
 1031 Tibet–Qinghai Plateau. Geological Publishing House, Beijing, pp. 1-125 (in Chinese).

1032 Lu, S.N., Yuan, G.B., 2003. Geochronology of early Precambrian magmatic activities
 1033 in Aketasdhtage, East Altyn Tagh. *Acta Geological Sinica* 77, 61–68 (in Chinese with
 1034 English abstract).

1035 Lu, S.N., Li, H.K., Zhang, C.L., Niu, G.H., 2008. Geological and geochronological
 1036 evidence for the Precambrian evolution of the Tarim Craton and surrounding
 1037 continental fragments. *Precambrian Research* 160, 94-107.

1038 Lu, H., Xiong, S., 2009. Magnetostratigraphy of the Dahonggou section, northern
 1039 Qaidam Basin and its bearing on Cenozoic tectonic evolution of the Qilian Shan and
 1040 Altyn Tagh Fault. *Earth and Planetary Science Letters* 288, 539-550.

1041 Lu, H., Wang, E., Meng, K., 2014. Paleomagnetism and anisotropy of magnetic
 1042 susceptibility of the Tertiary Janggalsay section (southeast Tarim basin): Implications
 1043 for Miocene tectonic evolution of the Altyn Tagh Range. *Tectonophysics* 618, 67-78.

1044 Ludwig, K.R., 2003. User's manual for Isoplot 3.0: a geochronological toolkit for
 1045 Microsoft Excel. Berkeley Geochronology Center. Spec. Pub., 4, pp. 1-71.

1046 Mao, L.G., Xiao, A.C., Wu, L., Li, B.L., Wang, L.Q., Lou, Q.Q., Dong, Y.P., Qin, S.H.,
 1047 2014. Cenozoic tectonic and sedimentary evolution of southern Qaidam Basin, NE
 1048 Tibetan Plateau and its implication for the rejuvenation of Eastern Kunlun Mountains.
 1049 *Science China Earth Sciences*, 1-14 (in Chinese).

1050 Meng, Q.R., Hu, J.M., Yang, F.Z., 2001. Timing and magnitude of displacement on
 1051 the Altyn Tagh fault: constraints from stratigraphic correlation of adjoining Tarim and
 1052 Qaidam basins, NW China. *Terra Nova* 13, 86-91.

1053 Meng, Q.R., Fang, X., 2008. Cenozoic tectonic development of the Qaidam Basin in
 1054 the northeastern Tibetan Plateau. *Geological Society of America Special Papers* 444,
 1055 1-24.

1056 Métivier, F., Gaudemer, Y., Tapponnier, P., Meyer, B., 1998. Northeastward growth of
 1057 the Tibet plateau deduced from balanced reconstruction of two depositional areas: The
 1058 Qaidam and Hexi Corridor basins, China. *Tectonics* 17, 823-842.

1059 Meyer, B., Tapponnier, P., Bourjot, L., Metivier, F., Gaudemer, Y., Peltzer, G.,

1060 Shunmin, G., Zhitai, C., 1998. Crustal thickening in Gansu - Qinghai, lithospheric
1061 mantle subduction, and oblique, strike-slip controlled growth of the Tibet plateau.
1062 *Geophysical Journal International* 135, 1-47.

1063 Molnar, P., Tapponnier, P., 1975. Cenozoic tectonics of Asia: Effects of a continental
1064 collision. *Science* 189, 419-426.

1065 Pei, J., Sun, Z., Wang, X., Zhao, Y., Ge, X., Guo, X., Li, H., Si, J., 2009. Evidence for
1066 Tibetan plateau uplift in Qaidam basin before Eocene-Oligocene boundary and its
1067 climatic implications. *Journal of Earth Science* 20, 430-437.

1068 Peltzer, G., Tapponnier, P., 1988. Formation and evolution of strike-slip faults, rifts,
1069 and basins during the India-Asia collision: An experimental approach. *Journal of*
1070 *Geophysical Research: Solid Earth* (1978–2012) 93, 15085-15117.

1071 QBGMR (Qinhai Bureau of Geology Mineral Resources), 1991. Regional Geology of
1072 Qinghai Province. Geological Publishing House, Beijing, pp. 1–662 (in Chinese).

1073 Qi, C., Li, X., Liang, X., Liu, Y., Tu, X., 2005. High-precision measurement of Hf
1074 isotopic reference values for the U-Pb geochronology standard zircons by
1075 multi-collector inductively coupled plasma-mass spectrometry. *Journal of Chinese*
1076 *Mass Spectrometry Society* 26, 149-154.

1077 Qiu, N.S., 2002. Tectono-thermal evolution of the Qaidam Basin, China: evidence
1078 from Ro and apatite fission track data. *Petroleum Geoscience* 8, 279-285.

1079 Ritts, B.D., Hanson, A.D., Zinniker, D., Moldowan, J.M., 1999. Lower-Middle
1080 Jurassic nonmarine source rocks and petroleum systems of the northern Qaidam basin,
1081 northwest China. *AAPG Bulletin* 83, 1980-2005.

1082 Ritts, B.D., Biffi, U., 2000. Magnitude of post-Middle Jurassic (Bajocian)
1083 displacement on the central Altyn Tagh fault system, northwest China. Geological
1084 Society of America Bulletin 112, 61-74.

1085 Ritts, B.D., Yue, Y., Graham, S.A., 2004. Oligocene-Miocene Tectonics and
1086 Sedimentation along the Altyn Tagh Fault, Northern Tibetan Plateau: Analysis of the
1087 Xorkol, Subei, and Aksay Basins. The Journal of geology 112, 207-229.

1088 Roger, F., Arnaud, N., Gilder, S., Tapponnier, P., Jolivet, M., Brunel, M., Malavieille,
1089 J., Xu, Z., Yang, J., 2003. Geochronological and geochemical constraints on Mesozoic
1090 suturing in east central Tibet. Tectonics 22, 1037.

1091 Roger, F., Jolivet, M., Malavieille, J., 2008. Tectonic evolution of the Triassic fold
1092 belts of Tibet. Comptes Rendus Geoscience 340, 180-189.

1093 Roger, F., Jolivet, M., Malavieille, J., 2010. The tectonic evolution of the
1094 Songpan-Garzê (North Tibet) and adjacent areas from Proterozoic to Present: A
1095 synthesis. Journal of Asian Earth Sciences 39, 254-269.

1096 Robinson, D.M., Dupont-Nivet, G., Gehrels, G.E., Zhang, Y., 2003. The Tula uplift,
1097 northwestern China: Evidence for regional tectonism of the northern Tibetan Plateau
1098 during late Mesozoic–early Cenozoic time. Geological Society of America Bulletin
1099 115, 35-47.

1100 Searle, M., 1996. Geological evidence against large-scale pre-Holocene offsets along
1101 the Karakoram Fault: Implications for the limited extrusion of the Tibetan plateau.
1102 Tectonics 15, 171-186.

1103 Searle, M., Elliott, J., Phillips, R., Chung, S. L., 2011. Crustal–lithospheric structure

1104 and continental extrusion of Tibet. *Journal of the Geological Society* 168, 633-672.

1105 Shang Erjie, 2001. Basic structural style of tertiary in the western part of the northern
1106 area in the Qaidam basin and its petroleum significance. *Geoscience* 12, 422-424 (in
1107 Chinese with English abstract).

1108 Shi, R.D., Yang, J.S., Wu, C.L., Tsuyoshi, I., Takafumi, H., 2006. Island arc volcanic
1109 rocks in the north Qaidam UHP belt, northern Tibet plateau: Evidence for
1110 ocean–continent subduction preceding continent–continent Subduction. *Journal of*
1111 *Asian Earth Sciences* 28, 151-159.

1112 Sobel, E.R., Arnaud, N., 1999. A possible middle Paleozoic suture in the Altyn Tagh,
1113 NW China. *Tectonics* 18, 64-74.

1114 Sobel, E.R., Arnaud, N., Jolivet, M., Ritts, B.D., Brunel, M., 2001. Jurassic to
1115 Cenozoic exhumation history of the Altyn Tagh range, northwest China, constrained
1116 by $^{40}\text{Ar}/^{39}\text{Ar}$ and apatite fission track thermochronology. *Geological Society of*
1117 *America Memoirs* 194, 247-267.

1118 Song, T., Wang, X., 1993. Structural styles and stratigraphic patterns of
1119 syndepositional faults in a contractional setting: Examples from Qaidam basin,
1120 northwestern China. *AAPG bulletin* 77, 102-117.

1121 Song, S.G., Zhang, L.F., Niu, Y.L., Su, L., Song, B., Liu, D.Y., 2006. Evolution from
1122 oceanic subduction to continental collision: a case study from the Northern Tibetan
1123 Plateau based on geochemical and geochronological data. *Journal of Petrology* 47,
1124 435-455.

1125 Song, S., Niu, Y., Su, L., Zhang, C., Zhang, L., 2014. Continental orogenesis from

1126 ocean subduction, continent collision/subduction, to orogen collapse, and orogen
 1127 recycling: The example of the North Qaidam UHPM belt, NW China. *Earth-Science*
 1128 *Reviews* 129, 59-84.

1129 Staisch, L.M., Niemi, N.A., Hong, C., Clark, M.K., Rowley, D.B., Currie, B., 2014. A
 1130 Cretaceous - Eocene depositional age for the Fenghuoshan Group, Hoh Xil Basin:
 1131 Implications for the tectonic evolution of the northern Tibet Plateau. *Tectonics* 33,
 1132 281-301.

1133 Sun, Z.M., Yang, Z.Y., Pei, J.L., Ge, X.H., Wang, X.S., Yang, T.S., Li, W.M., Yuan,
 1134 S.H., 2005. Magnetostratigraphy of Paleogene sediments from northern Qaidam Basin,
 1135 China: implications for tectonic uplift and block rotation in northern Tibetan plateau.
 1136 *Earth and Planetary Science Letters* 237, 635-646.

1137 Sun, Z.C., Jing, M.C., Sun, N.D., Lu, Y.L., Cao, L., 2007. Discussion on boundary
 1138 between the upper and lower members of Xiaganchaigou Formation of Paleogene in
 1139 Well Kun-2, Qaidam Basin. *Journal of Palaeogeography* 9, 611–618 (in Chinese with
 1140 English abstract).

1141 Sun, Y., Chen, Z., Chen, B., Han F., Zhou, Y., Hao, R. and Li, S., 2014. Cenozoic
 1142 Uplift and Denudation of the EW-trending Range of Northern Altun Mountains:
 1143 Evidence from Apatite Fission Track Data. *Acta Geoscientica Sinica* 37, 67-75 (in
 1144 Chinese with English abstract).

1145 Tapponnier, P., Peltzer, G., Armijo, R., 1986. On the mechanics of the collision
 1146 between India and Asia. *Geological Society, London, Special Publications* 19,
 1147 113-157.

1148 Tapponnier, P., Xu, Z.Q., Roger, F., Meyer, B., Arnaud, N., Wittlinger, G., Yang, J.S.,
1149 2001. Oblique stepwise rise and growth of the Tibet Plateau. *Science* 294, 1671-1677.

1150 Thomas, W.A., 2011. Detrital-zircon geochronology and sedimentary provenance.
1151 *Lithosphere* 3, 304-308.

1152 Wang, E., 1997. Displacement and timing along the northern strand of the Altyn Tagh
1153 fault zone, northern Tibet. *Earth and Planetary Science Letters* 150, 55-64.

1154 Wang, Y., Zhang, X., Wang, E., Zhang, J., Li, Q., Sun, G., 2005. $^{40}\text{Ar}/^{39}\text{Ar}$
1155 thermochronological evidence for formation and Mesozoic evolution of the
1156 northern-central segment of the Altyn Tagh fault system in the northern Tibetan
1157 Plateau. *Geological Society of America Bulletin* 117, 1336-1346.

1158 Wang, E., Xu, F.Y., Zhou, J.X., Wan, J., Burchfiel, B.C., 2006a. Eastward migration of
1159 the Qaidam basin and its implications for Cenozoic evolution of the Altyn Tagh fault
1160 and associated river systems. *Geological Society of America Bulletin* 118, 349-365.

1161 Wang, C., Liu, L., Che, Z.C., Chen, D.L., Zhang, A.D., Luo, J.H., 2006b. U–Pb
1162 geochronology and tectonic setting of the granitic gneiss in Jianggaleisayi Eclogite
1163 Belt. *Geological Journal of China Universities* 12, 74-82 (in Chinese with English
1164 abstract).

1165 Wang, L., Xiao, A.C., Gong, Q.L., Liu, D., Wu, L., Zhou, S.P., Shen, Z.Y., Lou, Q.Q.,
1166 Sun, X.W., 2010. The unconformity in Miocene sequence of western Qaidam Basin
1167 and its tectonic significance. *Science China Earth Sciences* 53, 1126-1133 (in
1168 Chinese).

1169 Wang, C., Liu, L., Yang, W.Q., Zhu, X.H., Cao, Y.T., Kang, L., Chen, S.F., Li, R.S.,

1170 He, S.P., 2013. Provenance and ages of the Altyn Complex in Altyn Tagh:
 1171 implications for the early Neoproterozoic evolution of northwestern China.
 1172 *Precambrian Research* 230, 193-208.

1173 Wang, C., Dai, J., Zhao, X., Li, Y., Graham, S.A., He, D., Ran, B., Meng, J., 2014a.
 1174 Outward-growth of the Tibetan Plateau during the Cenozoic: A review.
 1175 *Tectonophysics* 621, 1-43.

1176 Wang, C., Liu, L., Xiao, P.X., Cao, Y.T., Yu, H.Y., Meert, J.G., Liang, W.T., 2014b.
 1177 Geochemical and geochronologic constraints for Paleozoic magmatism related to the
 1178 orogenic collapse in the Qimantagh–South Altyn region, northwestern China. *Lithos*
 1179 202–203, 1-20.

1180 Wiedenbeck, M., Alle, P., Corfu, F., Griffin, W., Meier, M., Oberli, F., Quadt, A.v.,
 1181 Roddick, J., Spiegel, W., 1995. Three natural zircon standards for U–Th–Pb, Lu–Hf,
 1182 trace element and REE analyses. *Geostandards Newsletter* 19, 1-23.

1183 Wittlinger, G., Tapponnier, P., Poupinet, G., Mei, J., Danian, S., Herquel, G., Masson,
 1184 F., 1998. Tomographic evidence for localized lithospheric shear along the Altyn Tagh
 1185 fault. *Science* 282, 74-76.

1186 Wu, C.L., Yang, J.S., Ireland, T., Wooden, J.L., Li, H.B., Wan, Y.S., Shi, R.D., 2001.
 1187 Zircon SHRIMP ages of Aolaoshan granite from the south margin of Qilianshan and
 1188 its geological significance. *Acta Petrologica Sinica* 17, 215–221 (in Chinese with
 1189 English abstract).

1190 Wu, C.L., Wooden, J.L., Robinson, P.T., Gao, Y.H., Wu, S.P., Chen, Q.L., Mazdab,
 1191 F.K., Mattinson, C., 2009. Geochemistry and zircon SHRIMP U–Pb dating of

1192 granitoids from the west segment of the North Qaidam. *Science China, Series D Earth*
 1193 *Science* 52, 1771–1790 (in Chinese).
 1194 Wu C. L., Gao Y. H., Lei M., Qin H. P., Liu C. H., Li M. Z., Frost. B. R., Wooden J. L.,
 1195 2014. Zircon SHRIMP U-Pb dating, Lu-Hf isotopic characteristics and petrogenesis of
 1196 the Palaeozoic granites in Mangya area, south Altun, NW China. *Acta Petrologica*
 1197 *Sinica* 30, 2297-2323 (in Chinese with English abstract).
 1198 Wu, L., Xiao, A., Wang, L., Shen, Z., Zhou, S., Chen, Y., Wang, L., Liu, D., Guan, J.,
 1199 2011. Late Jurassic–Early Cretaceous Northern Qaidam Basin, NW China:
 1200 Implications for the earliest Cretaceous intracontinental tectonism. *Cretaceous*
 1201 *Research* 32, 552-564.
 1202 Wu, L., Xiao, A.C., Yang, S.F., Wang, L.Q., Mao, L.G., Wang, L., Dong, Y.P., Xu, B.,
 1203 2012a. Two-stage evolution of the Altyn Tagh Fault during the Cenozoic: new insight
 1204 from provenance analysis of a geological section in NW Qaidam Basin, NW China.
 1205 *Terra Nova* 24, 387-395.
 1206 Wu, L., Xiao, A., Wang, L., Mao, L., Wang, L., Dong, Y., Xu, B., 2012b. EW-trending
 1207 uplifts along the southern side of the central segment of the Altyn Tagh Fault, NW
 1208 China: Insight into the rising mechanism of the Altyn Mountain during the Cenozoic.
 1209 *Science China Earth Sciences* 55, 926-939 (in Chinese).
 1210 Wu, L., Xiao, A., Ma, D., Li, H., Xu, B., Shen, Y., Mao, L., 2014. Cenozoic fault
 1211 systems in southwest Qaidam Basin, northeastern Tibetan Plateau: Geometry,
 1212 temporal development, and significance for hydrocarbon accumulation. *AAPG*
 1213 *bulletin* 98, 1213-1234.

1214 Xia, W.C., Zhang, N., Yuan, X.P., Fan, L.S., Zhang, B.S., 2001. Cenozoic Qaidam
 1215 basin, China: A stronger tectonic inversed, extensional rifted basin. AAPG bulletin 85,
 1216 715-736.

1217 Xiao, W., Windley, B.F., Yong, Y., Yan, Z., Yuan, C., Liu, C., Li, J., 2009. Early
 1218 Paleozoic to Devonian multiple-accretionary model for the Qilian Shan, NW China.
 1219 Journal of Asian Earth Sciences 35, 323-333.

1220 Xiao, Q., Shao, G., Liu-Zeng, J., Oskin, M.E., Zhang, J., Zhao, G., Wang, J., 2015.
 1221 Eastern termination of the Altyn Tagh Fault, western China: Constraints from a
 1222 magnetotelluric survey. Journal of Geophysical Research: Solid Earth 120,
 1223 2838-2858.

1224 XBGMR (Xinjiang Bureau of Geology and Mineral Resources), 1993. Regional
 1225 Geology of Xinjiang Uygur Autonomous Region. Geological Publishing House,
 1226 Beijing, pp. 1-841 (in Chinese with English abstract).

1227 Yang, F., Ma, Z.Q., Xu, T.C., Ye, S.J., 1992. A Tertiary paleomagnetic stratigraphic
 1228 profile in Qaidam basin. Acta Petrolei Sinica 13, 97-101 (in Chinese with English
 1229 abstract).

1230 Yang, J., Xu, Z., Song, S., Wu, C., Shi, R., Zhang, J., Wan, Y., Li, H., Jin, X., Jolivet,
 1231 M., 2000. Discovery of eclogite in Dulan, Qinghai Province and its significance for
 1232 studying the HP-UHP metamorphic belt along the central orogenic belt of China. Acta
 1233 Geologica Sinica 74, 156-168 (in Chinese with English abstract).

1234 Yang, J., Xu, Z., Zhang, J., Chu, C. Y., Zhang, R., Liou, J. G., 2001. Tectonic
 1235 significance of early Paleozoic high-pressure rocks in Altun-Qaidam-Qilian

1236 Mountains, northwest China. Geological Society of America Memoirs 194, 151-170.

1237 Yang, M.H., Song, J.J., 2002. Petrology of the Lenghu granite mass, northwestern
1238 Qaidam basin, China. Northwestern Geol. 35, 94-98 (in Chinese with English
1239 abstract).

1240 Yang, J.S., Wu, C.L., Zhang, J.X., Shi, R.D., Meng, F.C., Wooden, J.L., Yang, H.Y.,
1241 2006. Protolith of eclogites in the north Qaidam and Altun UHP terrane, NW China:
1242 Earlier oceanic crust? Journal of Asian Earth Sciences 28, 185-204.

1243 Yang, W., Jolivet, M., Dupont-Nivet, G., Guo, Z., Zhang, Z., Wu, C., 2013. Source to
1244 sink relations between the Tian Shan and Junggar Basin (northwest China) from Late
1245 Palaeozoic to Quaternary: evidence from detrital U-Pb zircon geochronology. Basin
1246 Research 25, 219-240.

1247 Yang, W., Jolivet, M., Dupont-Nivet, G., Guo, Z., 2014. Mesozoic - Cenozoic tectonic
1248 evolution of southwestern Tian Shan: Evidence from detrital zircon U/Pb and apatite
1249 fission track ages of the Ulugqat area, Northwest China. Gondwana Research 26,
1250 986-1008.

1251 Yin, A., Harrison, T.M., 2000. Geologic evolution of the Himalayan-Tibetan orogen.
1252 Annual Review of Earth and Planetary Sciences 28, 211-280.

1253 Yin, A., Rumelhart, P., Butler, R., Cowgill, E., Harrison, T., Foster, D., Ingersoll, R.,
1254 Qing, Z., Xian-Qiang, Z., Xiao-Feng, W., 2002. Tectonic history of the Altyn Tagh
1255 fault system in northern Tibet inferred from Cenozoic sedimentation. Geological
1256 Society of America Bulletin 114, 1257-1295.

1257 Yin, A., Dang, Y.Q., Zhang, M., McRivette, M.W., Burgess, W.P., Chen, X.H., 2007.

1258 Cenozoic tectonic evolution of Qaidam basin and its surrounding regions (part 2):
1259 Wedge tectonics in southern Qaidam basin and the Eastern Kunlun Range. Geological
1260 Society of America Special Papers 433, 369-390.

1261 Yin, A., Dang, Y.Q., Wang, L.C., Jiang, W.M., Zhou, S.P., Chen, X.H., Gehrels, G.E.,
1262 McRivette, M.W., 2008a. Cenozoic tectonic evolution of Qaidam basin and its
1263 surrounding regions (Part 1): The southern Qilian Shan-Nan Shan thrust belt and
1264 northern Qaidam basin. Geological Society of America Bulletin 120, 813-846.

1265 Yin, A., Dang, Y.Q., Zhang, M., Chen, X.H., McRivette, M.W., 2008b. Cenozoic
1266 tectonic evolution of the Qaidam basin and its surrounding regions (Part 3): Structural
1267 geology, sedimentation, and regional tectonic reconstruction. Geological Society of
1268 America Bulletin 120, 847-876.

1269 Yu, X.J., Fu, S.T., Guan, S.W., Huang, B., Cheng, F., Cheng, X., Zhang, T., Guo, Z.J.,
1270 2014. Paleomagnetism of Eocene and Miocene sediments from the Qaidam basin:
1271 Implication for no integral rotation since the Eocene and a rigid Qaidam block.
1272 Geochemistry, Geophysics, Geosystems 15, 2109-2127.

1273 Yuan, H.L., Gao, S., Liu, X.M., Li, H.M., Günther, D., Wu, F.Y., 2004. Accurate U-Pb
1274 age and trace element determinations of zircon by laser ablation - inductively coupled
1275 plasma-mass spectrometry. Geostandards and Geoanalytical Research 28, 353-370.

1276 Yuan, D.Y., Ge, W.P., Chen, Z.W., Li, C.Y., Wang, Z.C., Zhang, H.P., Zhang, P.Z.,
1277 Zheng, D.W., Zheng, W.J., Craddock, W.H., Dayem, K.E., Duvall, A.R., Hough, B.G.,
1278 Lease, R.O., Champagnac, J.-D., Burbank, D.W., Clark, M.K., Farley, K.A., Garzione,
1279 C.N., Kirby, E., Molnar, P., Roe, G.H., 2013. The growth of northeastern Tibet and its

1280 relevance to large-scale continental geodynamics: A review of recent studies.
 1281 Tectonics 32, 2013TC003348.

1282 Yue, Y.J., Liou, J.G., 1999. Two-stage evolution model for the Altyn Tagh fault, China.
 1283 Geology 27, 227-230.

1284 Yue, Y.J., Ritts, B.D., Graham, S.A., 2001. Initiation and long-term slip history of the
 1285 Altyn Tagh Fault. International Geology Review 43, 1087-1093.

1286 Yue, Y., Ritts, B.D., Graham, S.A., Wooden, J.L., Gehrels, G.E., Zhang, Z., 2004a.
 1287 Slowing extrusion tectonics: lowered estimate of post-Early Miocene slip rate for the
 1288 Altyn Tagh fault. Earth and Planetary Science Letters 217, 111-122.

1289 Yue, Y.J., Ritts, B.D., Hanson, A.D., Graham, S.A., 2004b. Sedimentary evidence
 1290 against large strike-slip translation on the Northern Altyn Tagh fault, NW China. Earth
 1291 and Planetary Science Letters 228, 311-323.

1292 Yue, Y.J., Graham, S.A., Ritts, B.D., Wooden, J.L., 2005. Detrital zircon provenance
 1293 evidence for large-scale extrusion along the Altyn Tagh fault. Tectonophysics 406,
 1294 165-178.

1295 Zhang, J.X., Zhang, Z.M., Xu, Z.Q., Yang, J.S., Cui, J.W., 2001. Petrology and
 1296 geochronology of eclogites from the western segment of the Altyn Tagh, northwestern
 1297 China. Lithos 56, 187-206.

1298 Zhang, W., 2006. The High precise Cenozoic magnetostratigraphy of the Qaidam
 1299 basin and uplift of the northern Tibetan Plateau. (Doctoral dissertation) Lanzhou
 1300 University, Lanzhou, pp. 1–105 (in Chinese with English abstract).

1301 Zhang, J.X., Li, H.K., Meng, F.C., Xiang, Z.Q., Yu, S.Y., Li, J.P., 2011. Polyphase

1302 tectonothermal events recorded in metamorphic basement from the Altyn Tagh, the
 1303 southeastern margin of the Tarim basin, western China: constraint from U-Pb zircon
 1304 geochronology –Pb zircon geochronology. *Acta Petrologica Sinica* 27, 23-46.

1305 Zhang, Z.C., Guo, Z.J., Li, J.F., Tang, W.H., 2012. Mesozoic and Cenozoic
 1306 uplift-denudation along the Altyn Tagh fault, northwestern china: constraints from
 1307 apatite fission track data. *Quaternary Sciences* 32, 499-509 (in Chinese with English
 1308 abstract).

1309 Zhang, C.H., Cheng, F., Huang, G.P., Huang, Y.Q., Xing, C., Guan, B., Zhang, Q.Y,
 1310 and Xu, G., 2013a. Sediment and reservoir characteristics with reservoir evaluation of
 1311 the Lulehe Formation in Qie 16 block of Kunbei oilfield in Qaidam Basin. *Acta*
 1312 *Petrolei Sinica* 20, 2883–2894 (in Chinese with English abstract).

1313 Zhang, W., Fang, X., Song, C., Appel, E., Yan, M., Wang, Y., 2013b. Late Neogene
 1314 magnetostratigraphy in the western Qaidam Basin (NE Tibetan Plateau) and its
 1315 constraints on active tectonic uplift and progressive evolution of growth strata.
 1316 *Tectonophysics* 599, 107-116.

1317 Zhang, J., Mattinson, C., Yu, S., Li, Y., 2014a. Combined rutile–zircon thermometry
 1318 and U-Pb geochronology: New constraints on Early Paleozoic HP/UHT granulite in
 1319 the south Altyn Tagh, north Tibet, China. *Lithos* 200, 241-257.

1320 Zhang, H.P., Zhang, P.Z., Zheng, D.W., Zheng, W., Chen, Z.W., Wang, W.T., 2014b.
 1321 Transforming the Miocene Altyn Tagh fault slip into shortening of the north-western
 1322 Qilian Shan: insights from the drainage basin geometry. *Terra Nova* 26, 216-221.

1323 Zhao, J.M., Mooney, W.D., Zhang, X.K., Li, Z.C., Jin, Z.J., Okaya, N., 2006. Crustal

structure across the Altyn Tagh Range at the northern margin of the Tibetan plateau and tectonic implications. *Earth and Planetary Science Letters* 241, 804-814.

Zhou, J.X., Xu, F.Y., Wang, T.C., Cao, A.F., Yin, C.M., 2006. Cenozoic deformation history of the Qaidam Basin, NW China: Results from cross-section restoration and implications for Qinghai–Tibet Plateau tectonics. *Earth and Planetary Science Letters* 243, 195-210.

Zhuang, G., Hourigan, J.K., Ritts, B.D., Kent-Corson, M.L., 2011. Cenozoic multiple-phase tectonic evolution of the northern Tibetan Plateau: Constraints from sedimentary records from Qaidam basin, Hexi Corridor, and Subei basin, northwest China. *American Journal of Science* 311, 116-152.

Figure Caption

Fig. 1 (A) SRTM based digital topographic map of the Tibetan plateau. (B) Digital elevation model (DEM) and major tectonic elements of the Altyn Tagh Range, the Qaidam basin and the surrounding regions. The location of Fig. 2 is identified by the solid box. The DEM map was generated from the 90 m SRTM data. Note that the yellow solid line refers to the location of seismic profile AA'. (C) Interpreted and (D) non-interpreted seismic profile AA'. Note that the succession of Cenozoic depo-centers is marked along the long axis of the basin. These depo-centers gradually migrated eastward since the Eocene.

Fig. 2 Simplified geological map of the northwestern Qaidam basin and eastern

segment of the Altyn Tagh Range, adapted from the Geologic Map of the Tibetan plateau and adjacent areas compiled by the Chengdu Institute of Geology and Mineral Resources and Chinese Geological Survey (map scale, 1:1,500,000).

Fig. 3 Generalized stratigraphic column of the studied Mesozoic to Cenozoic series. Paleocurrent directions in the Huatugou section were obtained from Meng and Fang, (2008), Wu et al. (2012b) and Zhuang, et al. (2011). Paleocurrent directions in the Eboliang section were obtained from Zhuang, et al. (2011), Wu et al. (2012b) and Heermance et al. (2013). Note that sample CSL3 is cited from Cheng et al. (2015a), and samples HTG-E, HTG-N, CSL4, SZG1 and CSL5 are cited from Cheng et al. (2016). Red arrows refer to the general paleocurrent directions.

Fig. 4 Typical photomicrographs and field photographs of the various analyzed sediments. (A) Drill core sample of early to middle Jurassic (J_{1+2}) lithic sandstone obtained from drill well in the Eboliang section; (B) Thin section image of lithic sandstone from the Paleocene to early Eocene strata (E_{1+2}) in the Eboliang section, under crossed polarized light. (C) Horizontally stratified sandstone interbedded with mudstones of the Xiaganchaigou Formation (E_3^1) in the Eboliang section. (D) Thin section image of siltstone from the Oligocene strata (N_1), under crossed polarized light, obtained from drill well in the Eboliang section. (E) Thick-bedded sandstone in the early Miocene strata, Xiayoushashan Formation (N_2^1), Eboliang section. (F) Thin section image of siltstone from the Pliocene strata (N_2^3), in the Eboliang section,

under crossed polarized light. (G) Greenish white sandstone intercalated with mudstone in the middle Eocene strata (E_3^1), Huatugou section. (H) Greenish white sandstone intercalated with brownish red mudstone in the late Eocene strata (E_3^1), Huatugou section. (I) Thin section image of sandstone in the Oligocene strata (N_1), Huatugou section, under crossed polarized light.

Fig. 5 Representative CL images of zircons from the 11 sandstone samples. The yellow circles show the location of the U-Pb analysis. Numbers are U-Pb ages in Ma.

Fig. 6 U-Pb concordia diagrams for zircon grains of the 11 sandstone samples.

Fig. 7 Combined probability density functions (lines) and histogram plots (bars) depicting detrital zircon U-Pb ages of samples from the Eboliang section, arranged in stratigraphic order. Age distributions are colored according to age groups, and the pie diagrams show percentages of grains in those age categories. Paleocurrents rose diagrams in black are compiled from Heermance et al. (2013) and rose diagrams in gray are compiled from Wu et al. (2012b) and Ritts and Biffi (2000).

Fig. 8 Combined probability density functions (lines) and histogram plots (bars) depicting detrital zircon U-Pb ages of samples from the Huatugou section, arranged in stratigraphic order. Two distinctive age groups, indicative of separate sources, are colored. Paleocurrents rose diagrams are compiled from Meng and Fang, (2008), Wu

et al. (2012b) and Zhuang, et al. (2011).

Fig. 9 (A) Sketched geological map of the Altyn Tagh Range and surrounding areas, showing the distribution of zircon U–Pb ages for granitoids. The numbers denote ages in Ma. Relative probability plots of zircon U-Pb ages from basement rocks and intrusives in: (B) the Altyn Tagh Range, (C) the Qilian Shan, and (D) the Eastern Kunlun Range. Age data are mainly cited from: Guo and Li, (1999), Sobel and Arnaud, (1999), Yang et al. (2001), Zhang et al. (2001), Yang and Song, (2002), Cowgill et al. (2003), Gehrels et al. (2003a, 2003b), Jolivet et al. (2003), Lu and Yuan, (2003), Robinson et al. 2003, Roger et al. (2003), Chen et al. (2004), IGSQP, (2004), Yue et al. (2004, 2005), Liu et al. (2006), Shi et al. (2006), Song et al. (2006), Wang et al. (2006b), Yang et al. (2006), Lu et al. (2008), Roger et al. (2008), Bovet et al. (2009), Liu et al. (2009), Wu et al. (2009), Xiao et al. (2009), Roger et al. (2010), Zhang et al. (2011), Dai et al. (2013), Li et al. (2013), Wang et al. (2013), Long et al. (2014), Song et al (2014), Wang et al.(2014b) , Zhang et al. (2014a), Dong et al. (2014, 2015), Chen et al. (2015) and references therein. The blue arrow refers to the northeastward migration of the Huatugou and Eboliang sections from their origins. The red dashed line and the arrow show the ~80 km elongation of the early Neoproterozoic intrusions in the Altyn Tagh range.

Fig. 10 (A) Frequency histograms for restricted early Neoproterozoic ages groups of detrital zircons from the Paleocene to early Eocene Lulehe Fm. (E_{1+2}) and the Th/U

ratio for those zircon ages. (B) CL images of zircons for all early Neoproterozoic ages
zircons from the Paleocene to early Eocene Lulehe Fm. The yellow circles show the
location of the U-Pb analysis. Numbers are U-Pb ages in Ma. Note that the ~911 Ma
peak age for the Neoproterozoic ages groups coincides with the mean crystallization
age of ~910 Ma for the basement rocks in the western segment of the Altyn Tagh
Range (Fig. 9A; Wang et al., 2013). The distinctive Th/U ratio and oscillatory zones
for those zircons indicate magmatic sources.

Fig. 11 Seismic profiles in the southwestern Qaidam basin. See Fig. 2A for location.
Note the fold and thrusts in seismic profile BB', indicating the deformation in the
northwestern Qaidam basin since the early Miocene. Offset of the Paleogene strata in
seismic profiles CC' and DD' indicates the Eocene to Oligocene deformation in the
northwestern Qaidam basin which would be associated with deformation along the
ATF. Growth strata of the post-Oligocene strata in seismic profiles BB' and CC'
indicate intense tectonic movements along the ATF since the Miocene.

Fig. 12 Cenozoic kinematic model of the Altyn Tagh Fault and source to sink relation
between the western Qaidam basin and the surrounding regions. (A) during the
Paleocene to early Eocene, the basement rocks in the Altyn Tagh Range served as the
major source for the clastic material deposited in the Eboliang section, while
sediments in the Huatugou section were derived from the Eastern Kunlun Range; (B)
during the early Eocene to the Oligocene, the Qaidam basin migrated northeastward

due to left-lateral strike-slip faulting along the ATF; (3) since the Miocene, intense left-lateral faulting along the ATF continuously offset the Qaidam basin towards the northeast and triggered post-Oligocene crustal deformation within the Altyn Tagh Range and western Qaidam basin. See text for detailed discussion.

Table 1 Summary of the major characteristics and corresponding U-Pb age data for each sample.